



The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



Special article

Revisiting the Rich's formula: an update about granulomas in human tuberculosis

João Carlos Coelho Filho^{a,*}, Iukary Takenami^b, Sérgio Arruda^b

^a Laboratório de Patologia, Fundação José Silveira, Salvador, BA, Brazil

^b Laboratório Avançado de Saúde Pública, Centro de Pesquisas Gonçalo Moniz, FIOCRUZ, Salvador, BA, Brazil

ARTICLE INFO

Article history:

Available online 5 March 2013

Keywords:

Rich's formula

Tuberculosis

Immunopathogenesis

Granuloma

Hypersensitivity

ABSTRACT

The formula proposed by Rich in 1951 explained the formation in a tuberculous lesion in a period that was unknown cellular functions, cytokines and other immunological aspects involved in granuloma formation by tuberculosis; its components are assembled conceptually to explain the pathogenic mechanisms involved in the granulomatous lesion in tuberculosis. In this manuscript, we report an update of Rich's formula based on the new and old concepts about pathogenic mechanisms involved in the granulomatous lesion in tuberculosis. Current knowledge allows us to conclude that the balance between the characteristics of the bacillus and host protective response is necessary to indicate the outcome of pathogenesis, infection or active disease and the necrosis degree of the tuberculosis lesion.

© 2013 Elsevier Editora Ltda. Este é um artigo Open Access sob a licença de CC BY-NC-ND

Introduction

Tuberculosis (TB) is the most frequent cause of granulomas and is the most prevalent infectious disease in Brazil, remaining one of the most important problems of public health in the world. In 2010, 71,000 cases were reported, with a mortality of 4800/year.¹ Brazil is the 19th in rank among countries with the majority of TB cases in the world. Bahia, the north-east State of Brazil, is ranked 3rd in number of cases, after São Paulo and Rio de Janeiro. TB incidence in Bahia is 39.5/100,000 inhabitants.¹⁻³

TB is a granulomatous disease characterized by a cell-mediated hypersensitivity reaction in response to components of the wall structure of *Mycobacterium tuberculosis* especially cord factor and lipoarabinomannan.⁴ Granuloma

was first described in 1679 by Sylvius, and etymologically comes from the original word tubercle.⁵ Nowadays, granulomas are much better understood based on the knowledge of cellular immunology (cell mediated hypersensitivity reaction) and its interaction with cytokines and chemokines.

In 1951, Rich proposed that the pathogenesis of tuberculous lesion was directly dependent of inoculum, bacillus number (N), bacilli virulence (V) and host hypersensitivity (H), by counterbalancing and decreasing the lesion size, the natural and acquired resistance of the host (R_{n+a}). This formula can be represented by⁶:

$$L = \frac{N + V + H}{R_{(n+a)}}$$

Rich's formula shows components that are assembled conceptually to explain the pathogenic mechanisms involved in

* Corresponding author at: Laboratório de Patologia, Fundação José Silveira, João Carlos Coelho Filho, Fundação José Silveira, Ladeira do Campo Santo, Federação, s/n, Salvador, BA 40210-320, Brazil.

E-mail address: patologia@fjs.org.br (J.C. Coelho Filho).

1413-8670 © 2013 Elsevier Editora Ltda. Este é um artigo Open Access sob a licença de CC BY-NC-ND

<http://dx.doi.org/10.1016/j.bjid.2013.01.006>

the granulomatous lesion in TB. Considering the TB pathogenesis in two different clinical forms, infection and disease, where in infection, the balance depends on the host natural resistance. In contrast, in disease, the balance depends on the bacilli properties such as virulence and hypersensitivity phenomenon. In both clinical forms, either infection or disease, granulomas are detected in the tissue.⁷⁻⁹ Both stages are taking place inside of the granuloma, where the components of Rich's formula occur. The ancient concept was that a granuloma was a cellular agglomerate walling of the bacilli; thus in Rich's formula this should be considered in the denominator. This can be observed in some host, in which the granuloma formation is impaired and therefore the bacillus growth is faster. In fact, in these almost 60 years since the proposition of Rich's formula, the science has advanced significantly in the knowledge of the properties of *M. tuberculosis*, as well as in understanding of host defense mechanisms. These new concepts have been incorporated into the above formula.

Initiation of infection and granuloma formation

After mycobacterial arrival in the tissues of the respiratory tract, an unspecified reaction occurs in which neutrophils try to phagocytize the bacillus. This initial phase is not well studied in humans.¹⁰ Three weeks later, there is a development within these tissues of a more specific reaction, when mononuclear cells reach the focus of the infectious lesion, modulating the granuloma formation.^{9,11,12} Mononuclear cells including monocytes/macrophages and lymphocytes, cells with their predefined morphologies and functions, participate in the recognition and presentation of components of *M. tuberculosis* to the immune system.¹¹⁻¹³ During and after mononuclear cells naïve starts cells differentiation into specific cell types such as macrophages, dendritic cells, CD4 and CD8 T helper.¹⁴

In the process of granuloma formation, many macrophages with large cytoplasm and organelles with high phagocytic and microbicidal functions were observed.¹⁵⁻¹⁷ These cells can fuse into multinucleated giant cells; however, these changes are not completely understood.^{12,17} Mature macrophages can undergo a transformation into epithelioid cells (epithelioid histiocytes), which interdigitated their membranes aggregating into other cells as the epithelial barrier.¹⁶ CD4 and CD8 T helper (Th) lymphocytes as well as epithelioid monocytes and giant multinucleated cells participate in this granuloma formation that is orchestrated by INF- γ produced by CD4 and TNF- α secreted by macrophages.^{8,11,12,15,18} Other cells, such as neutrophils, join this granuloma via a dynamic process.^{9,19} All these facts have dynamic morphology that is influenced by the balance between host resistance and the *M. tuberculosis* virulence gene expression.^{9,11}

Cellular apoptosis of caseous necrosis can also be detected within the centers of granulomas. Progress has been made in determining these mechanisms; however, much remains not completely understood.⁹ One interesting point that has been found regards the function of the granuloma. Although the main consensus is that the granuloma that surrounds the *M. tuberculosis* is doing so in an attempt to restrict it from spreading.²⁰ Other experiments in the zebrafish have shown

that granuloma also contribute to bacilli spreading. During the granuloma's formation the tissues are injured and necrosis develops. This mechanism of lesion development depends on the acquired resistance versus the hypersensitivity developed by the patient to the bacillus during the infection.⁹

Granulomas seen in tuberculous pleurisy allow to distinguish the cellular components comparing the cells present in pleura fluid.²¹ Fayyazi et al. showed a possible apoptosis of epithelioid cells and the presence of CD4 T lymphocytes in the proximity of the focus of the necrosis. Such findings could suggest a direct relation between the cells, their secretions and the central necrosis of granulomas, in addition to the epithelioid cell's apoptosis.²² The interaction of all these aspects and also the elements of the extracellular matrix⁹ are part of the final product, the granuloma with central caseous necrosis.^{23,24}

Bacilli virulence

The genome sequence of *M. tuberculosis* allowed to conduct studies about the role of unknown genes through functional genomics. The identification of genes present in hypervirulent strains of *M. tuberculosis*,²⁵ but absent in pathogenic strains such as *Mycobacterium bovis* BCG, allowed a better understanding of the pathogenesis of TB.

Currently, the virulence of *M. tuberculosis* is defined by region of difference (RD1) genes, which are absent in BCG strains and are present in all *M. tuberculosis* virulent strains. Several studies have demonstrated that deletion of RD1 from *M. tuberculosis* results in attenuation of virulence, suggesting a dominant role in immune responses.^{26,27} Thus, deletion mutants of virulent *M. tuberculosis* strains for RD1 resemble BCG in their infectivity and attenuation.²⁷

Furthermore, another group of genes have been associated with virulence of *M. tuberculosis* in animal model. It is suggested that the mammalian cell entry genes (mce), which encodes Mce protein, were originally identified in *M. tuberculosis*²⁸ and have been associated with survival within macrophages and an increase in virulence.^{29,30} Disruption of the mce gene of BCG results in a mutant that exhibits reduced invasiveness epithelial cells, impairing their ability to survive and multiply inside phagocytic cells.³¹

Inhibition of macrophage apoptosis by *M. tuberculosis* strain H37Rv has been proposed as another virulence mechanism.³² This hypothesis is supported by studies that show that bacilli can cause necrosis instead of apoptosis in infected macrophages to avoid innate host defense. The macrophage cell death pathway by necrosis could be an exit mechanism for *M. tuberculosis* to facilitate the spread of infection and contribute to the formation of necrotic lesions in TB.^{33,34} Indeed, infection of macrophages with the attenuated *M. tuberculosis* causes apoptosis, which limits bacterial replications and promotes T-cell cross priming by antigen-presenting cells.³⁵

Host hypersensitivity

There is a strong relationship among immunity and hypersensitivity in TB. T lymphocytes can produce distinct types of

immune responses associated with different cytokine profiles and have opposing effects on resistance and susceptibility.¹⁴ The result of this balance will determine the degree of lesion in infected individuals and TB disease. Latent infection is associated with protective cellular immune responses whereas advanced disease is associated with immune suppression and exacerbated hypersensitivity reaction, causing tissue damage. Thus, T lymphocytes may also mediate harmful tissue-destroying hypersensitivity reactions that cause progression of disease.³⁶

Natural and acquired resistance

Macrophages, dendritic cells and neutrophils principally mediate the natural host resistance to *M. tuberculosis*. The available data on the role of neutrophils are controversial. Neutrophils are the first cells to be mobilized and arrive within hours at the site of infection. They are not professional phagocytes such as macrophages, but contribute substantially to innate resistance to infection. Neutrophils can also form an extracellular fibril matrix known as NETs (neutrophil extracellular traps). These NETs were described recently.³⁷ They are formed by activated neutrophils and consist of a DNA backbone with embedded antimicrobial peptides and enzymes. Thus, NETs represent a distinct innate defense mechanism to control and eliminate microbial infections.³⁸ On the other hand, the poor ability of neutrophils to restrict mycobacterial growth compared to alveolar macrophages indicates that the prevalence of neutrophils in TB inflammation contributes to the development of a nonspecific cellular reaction, rather than protection of the host, and that neutrophils may play the role of a "Trojan horse" for *M. tuberculosis*.^{10,39} Indeed, interferon-inducible neutrophil-driven blood transcriptional signature has been described in patients with active TB, which correlates with lung radiographic disease severity and implicates neutrophils directly in the pathogenesis of active TB.⁴⁰

Molecular studies show that the human genetic are responsible for natural resistance to *M. tuberculosis*. There is consensus in the literature that humans exposed to *Mycobacterium* millennially were evolutionarily selected, providing relatively more resistant cohorts. One example of this is the infection of Native Americans when Europeans first came to North America where the Apaches tribe was rapidly infected by acute and deadly TB.⁴¹ Also, indigenous Brazilians show a high TB rate, which can be attributable to an intrinsic susceptibility of these individuals.⁴²

The resistance mechanism in TB involves a great number of cellular interactions (macrophage, T cell). Several different T-cell populations are required for the successful control of the pathogen. This dynamic interplay underlying protection is the reason for the long-term persistence of *M. tuberculosis*. From the cellular point of view the T lymphocytes are differentiated in the thymus. CD4 and CD8 T-cells activate macrophages against *M. tuberculosis*. These cells recognize specific parts of the bacillus and secrete cytokines, which activate macrophages.^{9,11-15,18} T cells also act, induced by components of the bacillus, producing defense mechanisms. By reacting with proteins, a state of hypersensitivity is created that can be measured by the tuberculin test. In this

context, cytokines play an important role in the pathogenesis of TB. The cytokines are weighted molecules of about 8–14 kDa whose production is a function of the components of *M. tuberculosis*. These cytokines attract macrophages that transform into epithelioid cells.^{8,9,14} IFN- γ , TNF- α and interleukin-(IL)-12 are the most fundamental cytokines, but there are currently about 40 classified cytokines and other chemokines that act as receptors involved in the cellular dynamic of granulomas.⁴³

The events underlying the structural reorganization of immune cells to form a stable granuloma or progress to a pathologic lesion are dependent on other components. It is related with some components of the structure of *M. tuberculosis*, including some parts of its genetic material. These elements can mobilize, recruit, and amplify T cells reaction through the influence of cytokines.⁴⁴ Glycolipids, components of the *M. tuberculosis* wall (sero-lipidic membrane), also involve the bacillus and DNA. They contain a 6 kDa protein, designed as an ESAT-6, that mobilizes the T cell and activates the production of TNF- α and IFN- γ . Both of these activations lead the macrophages to destroy the *Mycobacterium*.^{19,45} Following the above events, as well as the arrival of bacilli in the tissues (for example, pulmonary alveoli), the formation of a granuloma with central caseous necrosis is completed.

Directed by T lymphocytes, modified monocytes aggregate and transform into epithelioid cells. With this fusion, giant multinucleated cells (Langerhans cells) are created. The dynamic previously described is also influenced by TNF- α .⁴⁴ In addition to Langerhans cells, TNF- α also plays a role in the formation of caseous necrosis within the center of the granuloma.^{9,44} The anaerobic environment and the action of phagolysosomes within the granulomas inhibit the spread of bacilli. The hypersensitivity promotes the progression to necrosis, making the cells more sensitive to the components of the bacillus (for example, tuberculostearic acid). The final differentiation of the granuloma will depend on the interplay of all the factors cited in this paper. When there is high immunity and low hypersensitivity there will be a low level of necrosis and the granuloma will most likely scar under the action of miofibroblasts. If the hypersensitivity is dominant, the necrosis will spread and the lesion will grow with an increase in the proliferation of bacilli.⁴⁶

In conclusion, although Rich's formula was proposed in the 1950s, it still holds some relevance to the distinct components within the granulomatous lesions. Even after incorporation of current knowledge about pathogenesis of *M. tuberculosis*, the balance between the characteristics of the bacillus and the host protective response is necessary to indicate the outcome of infection.

Conflict of interest

The authors have no conflict of interest to declare.

REFERENCES

1. Brazilian Ministry of Health. Available: <http://portalsaude.gov.br/portal/arquivos/pdf/apres.padrao.pnct.2011pdf>; 2011 [accessed June 2012].

2. World Health Organization. WHO report 2011: global tuberculosis control. Available at http://www.who.int/tb/publications/global_report/en/index.html; 2011 [accessed March 2012].
3. Ministério da Saúde, Secretaria de Políticas de Saúde Brazil. DATASUS. Brasília, Brazil: Ministério da Saúde. Available at <http://dtr2004.saude.gov.br/sinanweb/tabnet/tabnet> [accessed March 2012].
4. Rajni, Rao N, Meena LS. Biosynthesis and virulent behavior of lipids produced by *Mycobacterium tuberculosis*: LAM and cord factor: an overview. *Biotechnol Res Int*. 2011;2011:274693.
5. Sylvius F. *Opera Medica* (A. Wolfgang, 1679).
6. Rich AR. *The pathogenesis of tuberculosis*. 2 ed. IL, USA: Charles Thomas; 1951.
7. Rosemberg J. Mecanismo imunitário da tuberculose, síntese e atualização. *Boletim de Pneumologia Sanitária*. 2001;9:35-59.
8. Walzl G, Ronacher K, Hanekom W, Scriba TJ, Zumla A. Immunological biomarkers of tuberculosis. *Nat Rev Immunol*. 2011;11:343-54 [Review].
9. Ramakrishnan L. Revisiting the role of the granuloma in tuberculosis. *Nat Rev Immunol*. 2012;12:352-66.
10. Lowe DM, Redford PS, Wilkinson RJ, O'Garra A, Martineau AR. Neutrophils in tuberculosis: friend or foe? *Trends Immunol*. 2012;33:14-25.
11. Brighenti S, Andersson J. Local immune responses in human tuberculosis: learning from the site of infection. *J Infect Dis*. 2012;205 Suppl. 2:S316-24.
12. Russell DG, Cardona PJ, Kim MJ, Allain S, Altare F. Foamy macrophages and the progression of the human tuberculosis granuloma. *Nat Immunol*. 2009;10:943-8.
13. Schwander S, Dheda K. Human lung immunity against *Mycobacterium tuberculosis*: insights into pathogenesis and protection. *Am J Respir Crit Care Med*. 2011;183:696-707.
14. Kaufmann SH. Protection against tuberculosis: cytokines, T cells, and macrophages. *Ann Rheum Dis*. 2002;61 Suppl. 2:ii54-8.
15. Flynn JL, Chan J, Lin PL. Macrophages and control of granulomatous inflammation in tuberculosis. *Mucosal Immunol*. 2011;4:271-8.
16. Adams DO. The structure of mononuclear phagocytes differentiating in vivo. I. Sequential fine and histologic studies of the effect of Bacillus Calmette-Guerin (BCG). *Am J Pathol*. 1974;76:17-48.
17. Bouley DM, Ghori N, Mercer KL, Falkow S, Ramakrishnan L. Dynamic nature of host-pathogen interactions in *Mycobacterium marinum* granulomas. *Infect Immun*. 2001;69:7820-31.
18. Feng CG, Bean AG, Hooi H, Briscoe H, Britton WJ. Increase in gamma interferon-secreting CD8(+), as well as CD4(+), T cells in lungs following aerosol infection with *Mycobacterium tuberculosis*. *Infect Immun*. 1999;67:3242-7.
19. Chegou NN, Essone PN, Loxton AG, et al. Potential of host markers produced by infection phase-dependent antigen-stimulated cells for the diagnosis of tuberculosis in a highly endemic area. *PLoS One*. 2012;7:e38501.
20. Davis JM, Ramakrishnan L. The role of the granuloma in expansion and dissemination of early tuberculous infection. *Cell*. 2009;136:37-49.
21. Santana A, Loureiro C, Barbosa Jr A, Arruda S. Morphological analyses of caseous necrosis correlated to CD68+ and CD8+ cells in biopsies of tuberculous pleurisy. *Int J Morphol*. 2009;27:193-200.
22. Fayyazi A, Eichmeyer B, Soruri A, et al. Apoptosis of macrophages and T cells in tuberculosis associated caseous necrosis. *J Pathol*. 2000;191:417-25.
23. Volkman HE, Pozos TC, Zheng J, Davis JM, Rawls JF, Ramakrishnan L. Tuberculous granuloma induction via interaction of a bacterial secreted protein with host epithelium. *Science*. 2010;327:466-9.
24. Bold TD, Ernst JD. Who benefits from granulomas, mycobacteria or host? *Cell*. 2009;136:17-9.
25. Cole ST, Brosch R, Parkhill J, et al. Deciphering the biology of *Mycobacterium tuberculosis* from the complete genome sequence. *Nature*. 1998;393:537-44.
26. Lewis KN, Liao R, Guinn KM, et al. Deletion of RD1 from *Mycobacterium tuberculosis* mimics bacille Calmette-Guerin attenuation. *J Infect Dis*. 2003;187:117-23.
27. Gao LY, Guo S, McLaughlin B, Morisaki H, Engel JN, Brown EJ. A mycobacterial virulence gene cluster extending RD1 is required for cytolysis, bacterial spreading and ESAT-6 secretion. *Mol Microbiol*. 2004;53:1677-93.
28. Arruda S, Bomfim G, Knights R, Huima-Byron T, Riley LW. Cloning of an *M. tuberculosis* DNA fragment associated with entry and survival inside cells. *Science*. 1993;261:1454-7.
29. Shimono N, Morici L, Casali N, et al. Hypervirulent mutant of *Mycobacterium tuberculosis* resulting from disruption of the *mce1* operon. *Proc Natl Acad Sci USA*. 2003;100:15918-23.
30. Gioffre A, Infante E, Aguilar D, et al. Mutation in *mce* operons attenuates *Mycobacterium tuberculosis* virulence. *Microbes Infect*. 2005;7:325-34.
31. Flesselles B, Anand NN, Remani J, Loosmore SM, Klein MH. Disruption of the mycobacterial cell entry gene of *Mycobacterium bovis* BCG results in a mutant that exhibits a reduced invasiveness for epithelial cells. *FEMS Microbiol Lett*. 1999;177:237-42.
32. Chen M, Gan H, Remold HG. A mechanism of virulence: virulent *Mycobacterium tuberculosis* strain H37Rv, but not attenuated H37Ra, causes significant mitochondrial inner membrane disruption in macrophages leading to necrosis. *J Immunol*. 2006;176:3707-16.
33. Divangahi M, Chen M, Gan H, et al. *Mycobacterium tuberculosis* evades macrophage defenses by inhibiting plasma membrane repair. *Nat Immunol*. 2009;10:899-906.
34. Behar SM, Divangahi M, Remold HG. Evasion of innate immunity by *Mycobacterium tuberculosis*: is death an exit strategy? *Nat Rev Microbiol*. 2010;8:668-74.
35. Winau F, Weber S, Sad S, et al. Apoptotic vesicles crossprime CD8 T cells and protect against tuberculosis. *Immunity*. 2006;24:105-17.
36. Dannenberg Jr AM. Delayed-type hypersensitivity and cell-mediated immunity in the pathogenesis of tuberculosis. *Immunol Today*. 1991;12:228-33 [Research Support, U.S. Gov't, P.H.S. Review].
37. Brinkmann V, Reichard U, Goosmann C, et al. Neutrophil extracellular traps kill bacteria. *Science*. 2004;303:1532-5 [Research Support, U.S. Gov't, P.H.S.].
38. Wartha F, Beiter K, Normark S, Henriques-Normark B. Neutrophil extracellular traps: casting the NET over pathogenesis. *Curr Opin Microbiol*. 2007;10:52-6.
39. Eruslanov EB, Lyadova IV, Kondratieva TK, et al. Neutrophil responses to *Mycobacterium tuberculosis* infection in genetically susceptible and resistant mice. *Infect Immun*. 2005;73:1744-53.
40. Berry MP, Graham CM, McNab FW, et al. An interferon-inducible neutrophil-driven blood transcriptional signature in human tuberculosis. *Nature*. 2010;466:973-7.
41. Ferguson RG. *Studies in tuberculosis*. Toronto: Univ Tor Press; 1955.
42. Barbosa T, Benigno A, Farre L, Prat JG, Barral-Netto M, Arruda S. Tuberculosis among Yanomami communities from Alto Rio Negro, State of Amazonas, Brazil, 1990 to 2002. *Enf Emerg*. 2008;10:138-43.

43. Zlotnik A, Yoshie O. Chemokines: a new classification system and their role in immunity. *Immunity*. 2000;12:121-7.
44. Roach DR, Bean AG, Demangel C, France MP, Briscoe H, Britton WJ. TNF regulates chemokine induction essential for cell recruitment, granuloma formation, and clearance of mycobacterial infection. *J Immunol*. 2002;168:4620-7.
45. Guo S, Xue R, Li Y, Wang SM, Ren L, Xu JJ. The CFP10/ESAT6 complex of *Mycobacterium tuberculosis* may function as a regulator of macrophage cell death at different stages of tuberculosis infection. *Med Hypotheses*. 2012;78:389-92.
46. Adams DO. The granulomatous inflammatory response. A review. *Am J Pathol*. 1976;84:164-92.