



The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



Review article

Prevalence of human immunodeficiency virus/hepatitis C virus co-infection in Brazil and associated factors: a review

Valdete Meurer Kuehlkamp^a, Fabiana Schuelter-Trevisol^{a,b,*}

^a Postgraduate Program in Health Sciences, Universidade do Sul de Santa Catarina (UNISUL), Tubarão, SC, Brazil

^b Centro de Pesquisas Clínicas, Hospital Nossa Senhora da Conceição, Tubarão, SC, Brazil

ARTICLE INFO

Article history:

Received 27 September 2012

Accepted 29 October 2012

Available online 13 May 2013

Keywords:

HIV

Hepatitis C

Co-infection

Brazil

ABSTRACT

The hepatitis C virus and human immunodeficiency virus share the same transmission routes, which makes co-infection an unfavorable condition for the natural history of both viral diseases. In this context, it should be highlighted that the knowledge of the extent of co-infection and associated risk factors is a vital tool for prevention and control over infectious diseases. The aim of this study was to review the literature, seeking to examine the prevalence of human immunodeficiency virus/hepatitis C virus co-infection reported in studies conducted in Brazil, and identify the main risk factors associated with co-infection.

The electronic search was conducted in the Medline, Lilacs and SciELO databases. The following keywords were used: human immunodeficiency virus and Hepatitis C or hepatitis C virus and Brazil. The search led to 376 articles, of which 69 were selected for data extraction. We excluded animal studies, reports or case series, review articles, letters to the editor, other types of hepatitis and those studies in which co-infected patients were intentionally selected for comparison to single infected individuals. As a result, 40 articles were reviewed. The majority of the population in these studies was male (71%) and young adults, with a mean age of 26.7 years. The prevalence of hepatitis C virus co-infection among individuals living with human immunodeficiency virus in the studies conducted in Brazil ranged from 3.3% (serum samples) to 82.4% (drug users), with an average of 20.3%. The findings reveal that the prevalence of human immunodeficiency virus/hepatitis C virus co-infection is highly variable, depending on the characteristics of the study population. Risk factors associated with human immunodeficiency virus/hepatitis C virus co-infection were injection drug use and blood transfusion.

© 2013 Elsevier Editora Ltda. Este é um artigo Open Access sob a licença de [CC BY-NC-ND](https://creativecommons.org/licenses/by-nc-nd/4.0/)

Introduction

The human immunodeficiency virus (HIV) and hepatitis C virus (HCV) share the same transmission routes, which causes

high rates of co-infection, and is a severe clinical condition for both viral diseases.¹

HIV is most commonly spread by sexual contact with an infected partner, through contact with infected blood, accidental sticks or vertical transmission.² HCV transmission

* Corresponding author at: Programa de Pós-Graduação em Ciências da Saúde, Av. José Acácio Moreira, 787, Dehon, Tubarão, SC, 88704-900, Brazil.

E-mail addresses: fastrevisol@gmail.com, fabiana.trevisol@unisul.br (F. Schuelter-Trevisol).

1413-8670 © 2013 Elsevier Editora Ltda. Este é um artigo Open Access sob a licença de [CC BY-NC-ND](https://creativecommons.org/licenses/by-nc-nd/4.0/)

<http://dx.doi.org/10.1016/j.bjid.2012.10.023>

occurs mostly through infected blood and blood products, but sexual, accidental and vertical transmission has been also evidenced.³

The HIV/HCV co-infection is a significant risk factor for liver fibrosis, since HIV seropositivity and low CD4+ count seems to accelerate this process.^{4,5} The progression of fibrosis in HCV/HIV co-infected patients occurs with high frequency during an average estimated time of three years.⁶ Similarly, HCV infection has been shown to be associated with faster progression to acquired immunodeficiency syndrome (AIDS).⁷

In Brazil, according to the Epidemiological Bulletin on Viral Hepatitis (2011), co-infection rates reach a percentage of 11.4% in HIV-seropositive patients.⁸ This percentage shows regional variation being dependent on the characteristics of the study population. However, this estimate is based on cases reported by the Information System for Notifiable Diseases (SINAN), and may not be representative of the general population, since only AIDS cases are reported, and there is a relative delay between case detection and publishing of reports.⁸

Therefore, this study is intended to review this issue, seeking to examine the prevalence of HIV/HCV co-infection encountered in studies conducted in Brazil, and identify the main risk factors for this co-infection.

Methods

This is a narrative review on HIV/HCV co-infection in Brazil. The electronic search was conducted in the Medline, Lilacs and SciELO databases. The following keywords were used: HIV [MeSH] and HCV [textline] or Hepatitis C [MeSH] and Brazil [textline] in English and the corresponding words in Portuguese. The references of the selected articles were examined for possible inclusion of additional articles.

We included original scientific articles that presented data on the prevalence of HIV/HCV co-infection in different populations, of both sexes, conducted in Brazil.

We excluded studies that addressed animal studies, reports or case series, review articles, letters to the editor, other types of hepatitis and those in which co-infected patients were intentionally selected for comparison to single-infected patients.

Duplicate articles were excluded. We started reading the titles, then the abstracts, and later the full texts. Exclusion criteria were applied in all stages, always by consensus between two readers.

The selected articles were analyzed to characterize the study: year of publication, location and period of data collection, type of study, study population and sampling (number of investigated subjects, mean age and gender of participants), prevalence rates of HIV, HCV and HIV/HCV co-infection, and risk factors associated with the presence of co-infection.

HIV-infected individuals were considered those who had positive anti-HIV serology. HCV-infected individuals were considered those who had positive anti-HCV serology. The co-infection rates were taken directly from the articles, and when this information was not provided, they were calculated by dividing the number of individuals who had serological markers for HIV and HCV by the total number of participants in the study.

To calculate the mean age and the prevalence of HIV/HCV co-infection, we used descriptive statistics to find the weighted mean, taking into account the sample size of each study.

Results

The search led to 376 articles, of which 69 were selected for data extraction. After applying the exclusion criteria, 40 articles remained (Fig. 1). This study period covered articles published between April 1999 and June 2012.

Of the 40 articles analyzed, 36 were cross-sectional studies (one multisite), three cohort studies (two historical cohorts) and a descriptive study.

In total, we studied 48,791 subjects; of these, 28,819 (59.0%) were pregnant women, 13,483 (27.6%) were subjects or blood samples from HIV-infected patients, 2780 (5.7%) were individuals under restriction of freedom, 550 (1.1%) drug users, 3159 (6.6%) individuals with other characteristics, such as alcoholics, homeless people, garbage collectors, or patients attending various outpatient clinics.

Demographic characteristics available in the selected studies showed that the majority of participants were men (71%).⁹⁻³⁴ Of the 40 selected articles, men predominated in the samples of 26, with only one³⁵ showing a predominance of women. The samples of seven studies comprised only women.³⁵⁻⁴¹ The mean age was 26.7 years among the papers that presented this information.^{9-18,20-23,25,26,28,30-37,40-43} Eight of the selected papers had no data regarding gender and age of the participants.^{38,39,44-49}

In 20 articles,^{9-12,14-17,19-21,23,25,26,34,43-47} the study population consisted entirely of HIV-infected individuals or serum samples, and showed HIV/HCV co-infection rates ranging from 3.3%¹¹ to 54.7%⁴⁷ (Table 1), with an average of 19.5%. Including the studies with pregnant women^{36,40} and injection drug users,²⁹ whose participants were all infected with HIV as well, the weighted mean of HIV/HCV co-infection was 20.3% among 13,894 participants.

Five papers studied pregnant women, aiming to determine the prevalence of certain infectious diseases and/or verify the risk of vertical transmission.³⁶⁻⁴⁰ In these studies, the prevalence of HIV/HCV co-infection ranged from 0.00007³⁷ to 27.6%³⁹ (Table 2).

Among drug users^{18,29,35} the prevalence of HIV/HCV co-infection ranged from 15.1%³⁵ to 82.4%²⁹ (Table 3). Five studies assessing individuals in restraint of freedom^{24,28,32,33,48} found a prevalence of co-infection ranging from 1.6%²⁸ to 7.1%³² (Table 4).

The HIV/HCV co-infection rates ranged from 0.6%³⁰ among homeless people who used hostels to 82.4%²⁹ among injection drug users. Studies that have determined the prevalence of HIV/HCV co-infection in other populations are presented in Table 5.

Out of the 40 papers reviewed only 14 reported data on risk factors associated with HIV/HCV co-infection, and these highlighted the injection drug use as the main transmission mechanism,^{12,14,15,18,22,23,25,34,42,46} followed by blood transfusion.^{9,17,23,25} Behavioral risks such as tattooing,^{11,23} piercing¹¹ and exposure to infected blood,¹⁴ although to a

Table 1 – Description of the studies on HIV/HCV co-infection among HIV-infected patients.

Reference, year of publication	Study type and sample size	Target population, setting and study period	Gender/sex, age (mean ± SD)	Prevalence of HIV/HCV co-infection HIV/HCV n (%)	Risk factor associated with HIV/HCV co-infection
Carvalho FH et al., 2009 ⁹	Cross-sectional n = 345	Individuals living with HIV or AIDS attending the University Hospital of Recife (PE), from March to December 2003.	65% men	14 (4.1)	Blood transfusion
Corvino SM et al., 2007 ⁴⁴	Cross-sectional n = 150	HIV-infected patients monitored routinely at the Tropical Diseases and Diagnostic Imaging clinics, and the Hepato-Hemocentro of the Faculty of Medicine of Botucatu. ^a	NA	22 (14.7)	NA
Hadlich E et al., 2007 ¹⁰	Cross-sectional n = 61	HIV seropositive patients treated at a hospital in Porto Alegre (RS), between the year 2003 and September 2004.	59% men 38.0 ± 11.1 years	6 (10.0)	NA
Santos EO et al., 2008 ¹¹	Cross-sectional n = 299	Blood samples from HIV-infected patients treated at the University Hospital of Maceió (AL), from May to November 2005.	65.6% men 40.0 years	10 (3.3)	History of hepatitis/icterus and tattooing/piercing
Mendes-Correa MC et al., 2001 ¹²	Cross-sectional n = 1457	HIV-infected individuals attending the outpatient care center for patients with HIV/AIDS at the University of São Paulo (SP), from January to December 1996.	82.2% men 34.08 ± 7.19 years	258 (17.7)	Drug use and sexual transmission
Segurado AC et al., 2004 ¹⁴	Cohort n = 495	HIV-infected patients attending the Santos AIDS Reference Center, in Santos (SP), from February 1997 to January 1998.	55.4% men 35.7 years	179 (36.2)	IDU and blood exposure
Tovo CV et al., 2006 ¹⁵	Cross-sectional n = 330	Medical records of HIV-infected patients attending the Infectious Diseases Service, Hospital Nossa Senhora da Conceição, Porto Alegre (RS). ^a	62.4% men 34.4 ± 10.6 years	126 (38.2)	IDU
Victoria MB et al., 2010 ¹⁶	Descriptive n = 1582	AIDS patients reported to SINAN, who underwent serological testing for hepatitis C, Tropical Medicine Foundation of Amazonas, between 2000 and 2007.	72.9% men 38.5 ± 10.2 years	70 (4.42)	NA
Pavan MH, 2003 ¹⁷	Cross-sectional n = 221	Patients living with HIV in the city of Campinas (SP), between 1992 and 1995.	69.4% men 30.8 years	119 (53.8)	Blood transfusion
Wolf FH et al., 2010 ⁴⁵	Cross-sectional n = 1143	HIV patients under treatment at a referral outpatient center for HIV/AIDS of the Brazilian public health system, from March 2005 to September 2006.	NA	357 (31.2)	NA
Marins JR et al., 2005 ⁴⁶	Historic cohort n = 2821	Adult patients with AIDS in Brazil in 1995 and 1996. Data collected from medical records, between April 2000 and January 2002.	NA	833 (29.5)	IDU
Bonamigo RR et al., 2004 ¹⁹	Cross-sectional n = 105	HIV-infected patients with dermatoses treated at the Dermatology Clinic of the Department of Health of the State of Rio Grande do Sul. ^a	75.2% men	35 (38.0)	NA

Table 1 (Continued)

Reference, year of publication	Study type and sample size	Target population, setting and study period	Gender/sex, age (mean ± SD)	Prevalence of HIV/HCV co-infection HIV/HCV n (%)	Risk factor associated with HIV/HCV co-infection
Carmo RA et al., 2008 ²⁰	Historic cohort n = 824	HIV-infected patients treated at the Reference Center for Infectious Diseases, Belo Horizonte (MG), from January 1996 to 30 June 2001.	64.3% men	76 (9.2)	NA
Sampaio AS et al., 2009 ²¹	Cross-sectional n = 429	HIV-infected patients treated at the referral service to assist people with HIV infection at the Oswaldo Cruz University Hospital of Recife (PE), between July and September 2004.	60.1% men 39.3 years	46 (10.7)	NA
Mussi AD et al., 2007 ²³	Cross-sectional n = 1008	Individuals living with HIV treated at the referral center of Cuiabá (MT), between January 2004 and July 2004.	51.5% men 37.2 years	110 (10.9)	IDU, tattooing, blood transfusion before 1991.
Treitingner A et al., 1999 ⁴⁷	Cross-sectional n = 95	Blood samples from HIV-positive patients attending the Hospital Nereu Ramos or Health Centre II, from April 1994 to March 1995.	NA	52 (54.7)	
Monteiro MR et al., 2004 ²⁵	Cross-sectional n = 406	Patients living with HIV treated at the public health service in the city of Belém (PA), from September 1999 to April 2000.	74.1% men 34.2 ± 8.26 years	65 (16.0)	IDU, blood transfusion or blood products, and age > 49 years.
Morimoto HK et al., 2005 ²⁶	Cross-sectional n = 758	HIV-infected population attending various regional centers of the health referral center for AIDS in Paraná, from September 2001 to December 2002.	55.9% men 36.7 ± 9.5 years	159 (21.0)	NA
Braga EL et al., 2006 ²⁷	Cross-sectional n = 704	HIV patients attending the referral centers in the state of Amazonas, 2003.	65% men	35 (5.0)	Use of illicit drugs.
Santos KF et al., 2010 ³⁴	Cross-sectional n = 250	Medical records of HIV-positive patients monitored at the Clinic of Infectious Diseases, University Hospital of Santa Maria (RS), from June to November 2006.	76.9% men 33.8 ± 7.4 years	78 (31.2)	IDU

NA, not assessed; IDU, injection drug user; HIV, human immunodeficiency virus; HCV, hepatitis C virus; SD, standard deviation; n, sample size.

^a Study period not reported.

lesser extent, as well as sexual exposure,¹² homosexuality,¹⁸ and older age^{22,25} were positively associated with the presence of HIV/HCV co-infection.

Discussion

This review comprised 40 studies that surveyed a total of 48,791 Brazilians, including pregnant women, subjects or blood samples from HIV-infected patients, individuals under restriction of freedom, drug users, individuals with other characteristics, such as alcoholics, homeless people, garbage collectors, or patients attending various outpatient clinics. Due to this regional and population diversity, the prevalence of HIV/HCV co-infection had great amplitude. This variation

is explained by the exposure to different risk factors for co-infection.²

Thus, we chose to group the participants with similar characteristics, creating subgroups to analyze the seroprevalence rates of HIV/HCV co-infection.

Of the 20 studies that had determined the prevalence of HCV co-infection among individuals living with HIV, the average co-infection rate was 19.5%, which is higher than the rate of 11.8% published by the Ministry of Health in the last Epidemiological Bulletin of Viral Hepatitis.⁸ This can be attributed to the fact that this estimate is based on cases reported by the SINAN, and may not be representative of the general population, since only AIDS cases are reported and there is a delay between case detection and publishing of reports.⁸




Fig. 1 – Flowchart of the selection of articles.

The HIV/HCV co-infection rates among individuals in restraint of freedom are higher than in the general population. Containment is considered a risk factor due to crowding into small enclosures, promiscuity and rape, illicit drug use, and sharing of personal hygiene objects. These factors create favorable conditions for infection with HIV, HCV and co-infection.⁵⁰

According to the demographic characteristics of the participants in the different studies, 71% were men, which is in accordance with the national and international data showing that men are the most affected group by HIV, HCV and, consequently, co-infection.^{8,51,52} The mean age was 26.7 years, i.e., young adults, unlike some studies that showed older age populations.^{7,15} This discrepancy can be explained by the inclusion of teenagers³³ and pregnant women,³⁶⁻⁴⁰ e.g., women of childbearing age, and therefore, younger, coupled with the fact that several studies did not provide the age of the study population,^{19,38,39,44-49} which may have underestimated this figure.^{19,38,39,44-49} It should be emphasized, however, that the predominance of men and young population represents the sample described in the studies, and not the co-infection cases, since these data were not available. Nevertheless, one

can assume that there was a demographic similarity among the detected co-infection cases.

The articles that addressed studies with pregnant women showed a great variation of co-infection rates, ranging between 0.00007 and 27.6%.³⁶⁻⁴⁰ This variation can be explained by the fact that only Costa et al.³⁷ conducted research with large numbers of pregnant women to determine the prevalence of infection with HIV, HCV and co-infection, reaching rate values of 0.13%, 0.22% and 0.00007%, respectively. These findings diverge from some different studies included in the review, whose main objective was to verify the vertical transmission by studying subgroups of HIV- and HCV-seropositive subjects or both, thereby increasing the rates found. In Brazil, the prevalence of HIV infection in pregnant women is estimated to be 0.41%.⁵³ There is no national estimate of the prevalence of HCV infection among pregnant women, but the rate of HCV vertical transmission is 0.3%.⁸

Injection drug use is a major mode of HIV and HCV transmission.⁵⁴ In all studies included in this review, injection drug users had high rates of co-infection. This high rate of transmission can be attributed to the sharing of drug

Table 2 – Description of studies on HIV/HCV co-infection among pregnant women.

Reference, year of publication	Study type and sample size	Target population, setting and study period	Age (mean ± SD)	Prevalence n (%)		
				HIV	HCV	HIV/HCV
Tornatore M et al., 2012 ³⁶	Cross-sectional n = 130	HIV-infected pregnant women attending the HIV/AIDS center at the University Hospital of the Federal University of Rio Grande do Sul, from July 2006 to December 2008.	26.2 ± 5.69 years	130 (100.0)	14 (10.8)	14 (10.8)
Costa ZB et al., 2009 ³⁷	Cross-sectional n = 28,561	Pregnant women attending antenatal care of the public health system in Goiânia (GO), between 2004 and 2005.	23.9 ± 5.6 years	38 (0.13)	65 (0.22)	2 (0.00007)
Gardenal RV et al., 2011 ³⁸	Cross-sectional n = 23	HCV-positive pregnant women enrolled in the System Information Program for Humanization of Prenatal and Birth (SISPRENATAL) used by the Health Secretary of the municipality of Campo Grande (MS), between 2002 and 2005.	NA	2 (8.7)	23 (100.0)	2 (8.7)
Peixoto MF et al., 2004 ³⁹	Cross-sectional n = 29	HCV-positive pregnant women attending antenatal care of the Hospital Nossa Senhora da Conceição of Porto Alegre (RS), from August 1998 to November 1999.	NA	8 (27.6)	29 (100.0)	8 (27.6)
Dal Fabbro MM et al., 2005 ⁴⁰	Cross-sectional n = 76	HIV-infected pregnant women treated at the Reference Department in Campo Grande (MS), from May 1996 to October 2001.	24.0 years	76 (100.0)	11 (14.5)	11 (14.5)

NA, not assessed; HIV, human immunodeficiency virus; HCV, hepatitis C virus; SD, standard deviation; n, sample size.

preparation equipment, facilitating HIV and HCV transmission through the exposure to contaminated blood.⁵⁵ This pathway is the main route of transmission for HCV. Drug injection practices also predispose to other risky behaviors such as promiscuity, lack of condom use, homosexual intercourse, among others.⁵⁶

Different HIV/HCV co-infection rates were observed among the study population that participated in the studies included in this review. The heterogeneity of the studies did neither allow summarizing the prevalence of coinfection in each subgroup nor an overall prevalence. Therefore, we opted to calculate just the weighted average of HIV/HCV co-infection rate

Table 3 – Description of studies on HIV/HCV co-infection among drug users.

Reference, year of publication	Study type and sample size	Target population, setting and study period	Gender/sex, age (mean ± SD)	Prevalence n (%)			Risk factor associated with HIV/HCV co-infection
				HIV	HCV	HIV/HCV	
Zocratto KB, 2006 ¹⁸	Cross-sectional multisite study n = 272	Injecting drug users in 5 Brazilian cities selected by the AJUDE Brasil-I project, starting 1998.	82.7% men 29.25 ± 8.4 years	25 (9.2)	28 (10.3)	116 (42.6)	IDU, male homosexual intercourse
von Diemen L et al., 2010 ³⁵	Cross-sectional n = 73	Female crack users in Porto Alegre (RS), from August to December 2004.	100% women 28.1 ± 7.6 years	27 (37.0)	18 (27.7) ^a	11 (15.1)	NA
Marchesini AM et al., 2007 ²⁹	Cross-sectional n = 205	Injection drug users infected with HIV under treatment at the STD/AIDS municipal health department in São Paulo (SP), from January to November 2003.	81% men 39 ± 6.1 years	205 (100.0)	169 (82.4)	169 (82.4)	NA

HIV, human immunodeficiency virus; HCV, hepatitis C virus; NA, not assessed; SD, standard deviation; n, sample size; IDU, injection drug users; STD, sexually transmitted diseases; AIDS, acquired immunodeficiency syndrome.

^a The HCV infection rate of 27.7% was calculated based on a sample of 65 participants who underwent anti-HCV testing.

Table 4 – Description of studies on HIV/HCV co-infection among individuals in restraint of freedom.

Reference, year of publication	Study type and sample size	Target population, setting and study period	Gender/sex, age (mean ± SD)	Prevalence (%)		
				HIV	HCV	HIV/HCV
Pompilio MA et al., 2011 ²⁴	Cross-sectional n = 686	Prisoners of Mato Grosso do Sul State, 2009.	64.6% men	34 (5.0)	33 (4.8)	11 (1.6)
Gabe C and Lara GM, 2008 ⁴⁸	Cross-sectional n = 76	Serum samples of female prison inmates in Rio Grande do Sul, stored in the Santa Helena Laboratory, Porto Alegre (RS). ^a	NA	7 (9.2)	11 (14.5)	2 (2.6)
Catalan-Soares BC et al., 2000 ²⁸	Cross-sectional n = 63	Prisoners of Manhuaçu (MG). ^a	100% men 30.2 years	2 (3.2)	4 (6.3)	1 (1.6)
Guimaraes T et al., 2001 ³²	Cross-sectional n = 756	Blood samples of inmates at the House of Detention in São Paulo (SP), between December 1993 and January 1994.	100% men 30.2 years	105 (13.7)	310 (41.0)	54 (7.1)
Zanetta DM et al., 1999 ³³	Cross-sectional n = 1199	Teenagers from two FEBEM centers in São Paulo (SP), from December 1994 to April 1995.	92% men 16.2 years	38 (3.2)	21 (1.8)	21 (1.8)

HIV, human immunodeficiency virus; HCV, hepatitis C virus; NA, not assessed; SD, standard deviation; n, sample size; FEBEM, Portuguese acronym for the State Foundation for the Well-Being of Minors.

^a Study period not reported.

Table 5 – Description of studies on HIV/HCV co-infection among different population groups.

Reference, year of publication	Study type and sample size	Target population, setting and study period	Gender/sex, age (mean ± SD)	Prevalence (%)			Risk factor associated with HIV/HCV co-infection
				HIV	HCV	HIV/HCV	
Oliveira ML et al., 2009 ¹³	Cross-sectional n = 131	Alcoholic patients in a consecutive order, between March 1998 and November 1999.	90.8% men 44.3 ± 10.8 years	4 (3.0)	5 (4.2)	2 (1.5)	NA
Pereira GA et al., 2006 ⁴²	Cross-sectional n = 592	People attending a Center for anonymous testing and counseling for HIV in Goiânia (GO). ^a	65% women	19 (3.2)	15 (2.5)	14 (2.4)	IDU
Reis NR et al., 2011 ²²	Cross-sectional n = 402	Patients under treatment for tuberculosis (TB) at the referral hospital for infectious diseases in Goiânia (GO), from April 2008 to March 2010.	71.9% men 44.1 years	111 (27.6)	30 (7.5)	20 (5.0)	Age, IDU and HIV infection
Portelinho Filho AM et al., 2009 ⁴⁹	Cross-sectional n = 1.228	Individuals attending the Clinic of Infectious Diseases at the Dr. Dominic Leonardo Cerávolo Regional University Hospital in Presidente Prudente (SP), from January 2000 to December 2006.	NA	303 (24.7)	173 (14.0)	49 (4.0)	NA
Berra JAP et al., 2006 ⁴¹	Cross-sectional n = 225	Serum samples from patients with infectious diseases at the Rio Claro Regional Laboratory of the Adolfo Lutz Institute (SP), between March 2003 and June 2004.	100% women	8 (3.6)	17 (7.5)	2 (0.9)	NA
Brito VO et al., 2007 ³⁰	Cross-sectional n = 330	Homeless people who use hostels in Sao Paulo (SP), from 2002 to 2003.	80.9% men 40.2 years	6 (1.8)	28 (8.5)	2 (0.6)	NA
Rozman MA et al., 2008 ³¹	Cross-sectional n = 251	Seroepidemiological survey of autonomous garbage collectors in the city of Santos (SP), June 2005.	84.9% men 42.4 years	22 (8.9)	31 (12.4)	7 (2.8)	NA

NA, not assessed; HIV, human immunodeficiency virus; HCV, hepatitis C virus; SD, standard deviation; n, sample size; IDU, injection drug user.

^a Study period not reported.

in people living with HIV. The variation of coinfection rates found in this study is justified by the peculiar characteristics of the diverse population groups that are exposed to different risk factors.

The studies that reported data on risk factors associated with HIV/HCV co-infection highlighted that drug injection practices were the main transmission mechanism,^{12,14,15,18,22,23,25,34,46,57} followed by blood transfusion.^{9,17,23,25} Since the risk factors for HIV and HCV have close connection, these can be characterized as behavioral and biological factors.⁵⁸

The limitations of this review include missing information in some articles that could allow a more detailed analysis of the characteristics of the study populations and the risk factors for co-infection.

Conclusion

The data show that the prevalence of HIV/HCV co-infection is highly variable, depending on the prevalence of HIV and HCV infection, and the exposure to various risk factors in the study populations. The prevalence of HIV/HCV co-infection among people living with HIV was 20.3%. Risk factors associated with HIV/HCV co-infection were injection drug use and blood transfusion.

Conflict of interest

All authors declare to have no conflict of interest.

REFERENCES

- Miranda AE, Mercon-de-Vargas PR, Viana MC. Sexual and reproductive health of female inmates in Brazil. *Rev Saude Publica*. 2004;38:255-60.
- Soriano V, Garcia-Samaniego J, Rodriguez-Rosado R, Gonzalez J, Pedreira J. Hepatitis C and HIV infection: biological, clinical, and therapeutic implications. *J Hepatol*. 1999;31 Suppl. 1:119-23.
- Yen T, Keeffe EB, Ahmed A. The epidemiology of hepatitis C virus infection. *J Clin Gastroenterol*. 2003;36:47-53.
- Ragni MV, Belle SH. Impact of human immunodeficiency virus infection on progression to end-stage liver disease in individuals with hemophilia and hepatitis C virus infection. *J Infect Dis*. 2001;183:1112-5.
- Benhamou Y, Bochet M, Di Martino V, et al. Liver fibrosis progression in human immunodeficiency virus and hepatitis C virus coinfecting patients. The Multivirc Group. *Hepatology*. 1999;30:1054-8.
- Macias J, Berenguer J, Japon MA, et al. Fast fibrosis progression between repeated liver biopsies in patients coinfecting with human immunodeficiency virus/hepatitis C virus. *Hepatology*. 2009;50:1056-63.
- Rockstroh JK, Mocroft A, Soriano V, et al. Influence of hepatitis C virus infection on HIV-1 disease progression and response to highly active antiretroviral therapy. *J Infect Dis*. 2005;192:992-1002.
- Brasil. Boletim Epidemiológico Hepatites Virais. Brasília: Ministério da Saúde; 2011.
- Carvalho FH, Coelho MR, Vilella Tde A, Silva JL, Melo HR. HIV/HCV coinfection at an university hospital in Recife. *Brazil Rev Saude Publica*. 2009;43:133-9.
- Hadlich E, Alvares-Da-Silva MR, Dal Molin RK, Zenker R, Goldani LZ. Hepatitis C virus (HCV) viremia in HIV-infected patients without HCV antibodies detectable by third-generation enzyme immunoassay. *J Gastroenterol Hepatol*. 2007;22:1506-9.
- Santos EO, Coelho MRCD, Vilella TAS, Silva JLA, Neto EPAL. Ocorrência e fatores de risco para co-infecção pelo vírus da hepatite C (HCV) em pacientes com o vírus da imunodeficiência humana (HIV) em Maceió. *Brasil Rev Para Med*. 2008;22:21-7.
- Mendes-Correa MC, Barone AA, Guastini C. Hepatitis C virus seroprevalence and risk factors among patients with HIV infection. *Rev Inst Med Trop Sao Paulo*. 2001;43:15-9.
- Oliveira ML, Yoshida CF, Telles PR, et al. Trends in HCV prevalence, risk factors and distribution of viral genotypes in injecting drug users: findings from two cross-sectional studies. *Epidemiol Infect*. 2009;137:970-9.
- Segurado AC, Braga P, Etzel A, Cardoso MR. Hepatitis C virus coinfection in a cohort of HIV-infected individuals from Santos, Brazil: seroprevalence and associated factors. *AIDS Patient Care STDS*. 2004;18:135-43.
- Tovo CV, Dos Santos DE, de Mattos AZ, de Almeida PR, de Mattos AA, Santos BR. Ambulatorial prevalence of hepatitis B and C markers in patients with human immunodeficiency virus infection in a general hospital. *Arq Gastroenterol*. 2006;43:73-6.
- Victoria MB, Victoria Fda S, Torres KL, Kashima S, Covas DT, Malheiro A. Epidemiology of HIV/HCV coinfection in patients cared for at the Tropical Medicine Foundation of Amazonas. *Braz J Infect Dis*. 2010;14:135-40.
- Pavan MH, Aoki FH, Monteiro DT, Goncales NS, Escanhoela CA, Goncales Junior FL. Viral hepatitis in patients infected with human immunodeficiency virus. *Braz J Infect Dis*. 2003;7:253-61.
- Zocchetto KB, Caiaffa WT, Proietti FA, Carneiro-Proietti AB, Mingoti SA, Ribeiro GJ. HCV and HIV infection and co-infection: injecting drug use and sexual behavior, AJUDE-Brasil I Project. *Cad Saude Publica*. 2006;22:839-48.
- Bonamigo RR, Borges K, Rietjens J, Arenzon S, Blanco LF, Loureiro R. Human T lymphotropic virus 1 and hepatitis C virus as risk factors for inflammatory dermatoses in HIV-positive patients. *Int J Dermatol*. 2004;43:568-70.
- Carmo RA, Guimaraes MD, Moura AS, et al. The influence of HCV coinfection on clinical, immunological and virological responses to HAART in HIV-patients. *Braz J Infect Dis*. 2008;12:173-9.
- Sampaio AS, Alencar LCA, Moura PMMF, Correia JB, Barreto SB, Castelo A. Prevalencia de la co-infecção con hepatitis B y C en pacientes HIV positivos y factores de riesgo asociados. *Actualizaciones em SIDA*. 2009;17:12-7.
- Reis NR, Lopes CL, Teles SA, et al. Hepatitis C virus infection in patients with tuberculosis in Central Brazil. *Int J Tuberc Lung Dis*. 2011;15:1397-402.
- Mussi AD, Pereira RA, Correa e Silva Vde A, Martins RM, Souto FJ. Epidemiological aspects of hepatitis C virus infection among HIV-infected individuals in Mato Grosso State, Central Brazil. *Acta Trop*. 2007;104:116-21.
- Pompilio MA, Pontes ERJC, Castro ARCM, et al. Prevalence and epidemiology of chronic hepatitis C among prisoners of Mato Grosso do Sul State, Brazil. *J Venom Anim Toxina Incl Trop Dis*. 2011;17:216-22.
- Monteiro MR, do Nascimento MM, Passos AD, Figueiredo JF. Hepatitis C: prevalence and risk factors among patients with HIV/AIDS in Belem Para, in Brazilian Amazon. *Rev Soc Bras Med Trop*. 2004;37 Suppl. 2:40-6.

26. Morimoto HK, Caterino-De-Araujo A, Morimoto AA, et al. Seroprevalence and risk factors for human T cell lymphotropic virus type 1 and 2 infection in human immunodeficiency virus-infected patients attending AIDS referral center health units in Londrina and other communities in Parana, Brazil. *AIDS Res Hum Retroviruses*. 2005;21:256-62.
27. Braga EL, Lyra AC, Ney-Oliveira F, et al. Clinical and epidemiological features of patients with chronic hepatitis C co-infected with HIV. *Braz J Infect Dis*. 2006;10:17-21.
28. Catalan-Soares BC, Almeida RT, Carneiro-Proietti AB. Prevalence of HIV-1/2, HTLV-I/II, hepatitis B virus (HBV), hepatitis C virus (HCV), *Trypanosoma cruzi* among prison inmates at Manhuacu, Minas Gerais State, Brazil. *Rev Soc Bras Med Trop*. 2000;33:27-30.
29. Marchesini AM, Pra-Baldi ZP, Mesquita F, Bueno R, Buchalla CM. Hepatitis B and C among injecting drug users living with HIV in Sao Paulo, Brazil. *Ver Saude Publica*. 2007;41 Suppl. 2:57-63.
30. Brito VO, Parra D, Facchini R, Buchalla CM. HIV infection, hepatitis B and C and syphilis in homeless people, in the city of Sao Paulo, Brazil. *Rev Saude Publica*. 2007;41 Suppl. 2:47-56.
31. Rozman MA, Alves IS, Porto MA, et al. HIV infection and related risk behaviors in a community of recyclable waste collectors of Santos, Brazil. *Rev Saude Publica*. 2008;42:838-43.
32. Guimaraes T, Granato CF, Varella D, Ferraz ML, Castelo A, Kallas EG. High prevalence of hepatitis C infection in a Brazilian prison: identification of risk factors for infection. *Braz J Infect Dis*. 2001;5:111-8.
33. Zanetta DM, Strazza L, Azevedo RS, et al. HIV infection and related risk behaviours in a disadvantaged youth institution of Sao Paulo, Brazil. *Int J STD AIDS*. 1999;10:98-104.
34. Santos KF, Vieira TB, Beck STB, Leal DBR. Alterações laboratoriais encontradas em indivíduos co-infectados pelo vírus da imunodeficiência humana (HIV) e pelo vírus da hepatite C (HCV). *RBAC*. 2010;42:21-4.
35. von Diemen L, De Boni R, Kessler F, Benzano D, Pechansky F. Risk behaviors for HCV- and HIV-seroprevalence among female crack users in Porto Alegre. *Brazil Arch Womens Ment Health*. 2010;13:185-91.
36. Tornatore M, Goncalves CV, Bianchi MS, et al. Co-infections associated with human immunodeficiency virus type 1 in pregnant women from southern Brazil: high rate of intraepithelial cervical lesions. *Mem Inst Oswaldo Cruz*. 2012;107:205-10.
37. Costa ZB, Machado GC, Avelino MM, et al. Prevalence and risk factors for Hepatitis C and HIV-1 infections among pregnant women in Central Brazil. *BMC Infect Dis*. 2009;9:116.
38. Gardenal RV, Figueiro-Filho EA, Luft JL, et al. Hepatitis C and pregnancy: an analysis of factors associated with vertical transmission. *Rev Soc Bras Med Trop*. 2011;44:43-7.
39. Peixoto MF, Mattos AA, Remiao JO, Alexandre CO, Lemos SK, Azevedo AM. Vertical transmission of hepatitis C virus in a hospital in southern Brazil. *Arq Gastroenterol*. 2004;41:84-7.
40. Dal Fabbro MM, Cunha RV, Paniago AM, Lindenberg Ade S, Freitas GM, Nogueira SA. Prospective study on the prevention of vertical transmission of HIV in Campo Grande, Mato Grosso do Sul, Brazil, from 1996 to 2001. *Braz J Infect Dis*. 2005;9:20-7.
41. Berra JAP, Bacetti LB, Buzo AA. Soroprevalência de HIV, sífilis, hepatite B e C em mulheres do Centro de Ressocialização Feminino, Rio Claro, São Paulo. *Rev Inst Adolfo Lutz*. 2006;65:133-6.
42. Pereira GA, Stefani MM, Martelli CM, et al. Human immunodeficiency virus type 1 and hepatitis C virus co-infection and viral subtypes at an HIV testing center in Brazil. *J Med Virol*. 2006;78:719-23.
43. Braga WS, da Costa Castilho M, dos Santos IC, Moura MA, Segurado AC. Low prevalence of hepatitis B virus, hepatitis D virus and hepatitis C virus among patients with human immunodeficiency virus or acquired immunodeficiency syndrome in the Brazilian Amazon basin. *Rev Soc Bras Med Trop*. 2006;39:519-22.
44. Corvino SM, Henriques RMS, Grotto RMT, Pardini MIMC. Co-infecção HIV/HCV em pacientes de Botucatu e região. *Rev Bras Epidemiol*. 2007;10:537-43.
45. Wolff FH, Fuchs SC, Barcellos NN, et al. Co-infection by hepatitis C virus in HIV-infected patients in southern Brazil: genotype distribution and clinical correlates. *PLoS ONE*. 2010;5:e10494.
46. Marins JR, Barros MB, Machado H, Chen S, Jamal LF, Hearst N. Characteristics and survival of AIDS patients with hepatitis C: the Brazilian National Cohort of 1995-1996. *AIDS*. 2005;19 Suppl. 4:S27-30.
47. Treitinger A, Spada C, Silva EL, et al. Prevalence of serologic markers of HBV and HCV infection in HIV-1 seropositive patients in Florianopolis, Brazil. *Braz J Infect Dis*. 1999;3:1-5.
48. Gabe C, Lara GM. Prevalência de anti-HCV, anti-HIV e co-infecção HCV/HIV em um presídio feminino do Estado do Rio Grande do Sul. *RBAC*. 2008;40:87-9.
49. Portelinha Filho AM, Nascimento CU, Tannouri TN, et al. Seroprevalence of HBV, HCV and HIV co-infection in selected individuals from state of Sao Paulo, Brazil. *Mem Inst Oswaldo Cruz*. 2009;104:960-3.
50. Gois SM, Santos Junior HP, Silveira Mde F, Gaudencio MM. Beyond bars and punishments: a systematic review of prison health. *Cien Saude Colet*. 2012;17:1235-46.
51. Loko MA, Salmon D, Carrieri P, et al. The French national prospective cohort of patients co-infected with HIV and HCV (ANRS CO13 HEPAVIH): early findings, 2006-2010. *BMC Infect Dis*. 2010;10:303.
52. Mboti CI, Fielder M, Davies-Russell A, Jewell AP. Prevalence of HIV-1, HIV-2, hepatitis C and co-infection in The Gambia. *West Afr J Med*. 2009;28:16-9.
53. Brasil. Boletim Epidemiológico Aids e DST. Brasília: Ministério da Saúde; 2010.
54. Napoli C, Tafuri S, Pignataro N, Tedesco G, Maria AS, Quarto M. Risk factors for HBV/HIV/HCV in drug addicts: a survey of attendees of a department of pathological dependence. *J Prev Med Hyg*. 2010;51:101-4.
55. Yan YX, Gao YQ, Sun X, et al. Prevalence of hepatitis C virus and hepatitis B virus infections in HIV-positive Chinese patients. *Epidemiol Infect*. 2011;139:354-60.
56. Tedder RS, Gilson RJ, Briggs M, et al. Hepatitis C virus: evidence for sexual transmission. *BMJ*. 1991;302:1299-302.
57. Vogler IH, Nishiya A, Morimoto HK, et al. Serological, epidemiological and molecular aspects of hepatitis C virus infection in a population from Londrina, PR, Brazil, 2001-2002. *Rev Inst Med Trop Sao Paulo*. 2004;46:303-8.
58. Zhang Y, Chen P, Lu R, et al. Prevalence of HIV among men who have sex with men in Chongqing, China, 2006-2009: cross-sectional biological and behavioural surveys. *Sex Transm Infect*. 2012;88:444-50.