


The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



Original article

Parainfluenza virus as a cause of acute respiratory infection in hospitalized children[☆]



Rogério Pecchini^a, Eitan Naaman Berezin^{a,*}, Maria Cândida Souza^b, Lourdes de Andrade Vaz-de-Lima^b, Neuza Sato^b, Maristela Salgado^b, Mirthes Ueda^b, Saulo Duarte Passos^c, Raphael Rangel^d, Ana Catebelota^d

^a Medical Shool, Santa Casa de São Paulo, São Paulo, SP, Brazil

^b Center for Immunology, Instituto Adolfo Lutz, São Paulo, SP, Brazil

^c Faculdade de Medicina de Jundiaí, Jundiaí, SP, Brazil

^d Irmandade da Santa Casa de Misericórdia de São Paulo, SP, Brazil

ARTICLE INFO

Article history:

Received 27 December 2014

Accepted 31 March 2015

Available online 25 April 2015

ABSTRACT

Background: Human parainfluenza viruses account for a significant proportion of lower respiratory tract infections in children.

Objective: To assess the prevalence of Human parainfluenza viruses as a cause of acute respiratory infection and to compare clinical data for this infection against those of the human respiratory syncytial virus.

Methods: A prospective study in children younger than five years with acute respiratory infection was conducted. Detection of respiratory viruses in nasopharyngeal aspirate samples was performed using the indirect immunofluorescence reaction. Length of hospital stay, age, clinical history and physical exam, clinical diagnoses, and evolution (admission to Intensive Care Unit or general ward, discharge or death) were assessed. Past personal (premature birth and cardiopathy) as well as family (smoking and atopy) medical factors were also assessed.

Results: A total of 585 patients were included with a median age of 7.9 months and median hospital stay of six days. No difference between the HRSV+ and HPIV+ groups was found in terms of age, gender or length of hospital stay. The HRSV+ group had more fever and cough. Need for admission to the Intensive Care Unit was similar for both groups but more deaths were recorded in the HPIV+ group. The occurrence of parainfluenza peaked during the autumn in the first two years of the study.

Conclusion: Parainfluenza was responsible for significant morbidity, proving to be the second-most prevalent viral agent in this population after respiratory syncytial virus. No difference in clinical presentation was found between the two groups, but mortality was higher in the HPIV+ group.

© 2015 Published by Elsevier Editora Ltda.

[☆] Approval Number in Internal Review Board (Research Ethical Committee) – 064/09.

* Corresponding author.

E-mail address: eberezin2003@yahoo.com (E.N. Berezin).

<http://dx.doi.org/10.1016/j.bjid.2015.03.002>

1413-8670/© 2015 Published by Elsevier Editora Ltda.

Introduction

Acute respiratory tract infections (ARTIs) in childhood are a major cause of morbidity and mortality, with pneumonia being the leading cause of death among children between one and five years old.^{1,2} Viruses are the predominant etiology of ARTIs in this pediatric group.^{3–5}

The human parainfluenza viruses (HPIVs) account for a significant proportion of lower respiratory tract infections in children, where their rates of detection vary with pathology (respiratory infection of the upper or lower tract) and with the investigation setting (outpatient units or hospital wards).⁶

With the attention of parents and medical professionals firmly focused on annual outbreaks of respiratory syncytial virus (RSV) and influenza, HPIV infections can be overlooked.⁷ However, HPIVs are the second most common cause of lower and upper respiratory tract infections in children younger than five years after RSV.^{8–11}

This prospective study was conducted to determine the prevalence of HPIVs in children hospitalized for ARTI, assess the clinical characteristics of HPIV+ and RSV+ patients, and to compare the two groups identifying any specific features of infections caused by HPIVs. Seasonality was also examined in the present study.

Methods

This prospective study included all children with ARTI younger than five years with prodromes of upper respiratory tract infections (URTIs) on clinical history, admitted to the hospital general wards or Intensive Care Unit (ICU) of the Department of Pediatrics of Santa Casa de São Paulo Hospital (São Paulo city), between February 2005 and May 2007. After 2007, surveillance continued but using other methodology for diagnosing viral infections. Immunosuppressed patients like HIV-infected, and those with neoplasia or transplantations were excluded.

The Central Hospital at Santa Casa de São Paulo is a reference university hospital for the central and northern region of the city providing high complexity care, including emergency care. Approximately 60,000 patients receive medical attention annually at urgency and emergency units. Out of 9500 admissions during the study period, around one-third was due to respiratory infections. Data on clinical characteristics and demographics were collected at patient admission. The study was approved by the Research Ethics Committee of the Hospital under the number 064/09. Prior to inclusion in the study, parents or legal guardians of the patients signed a free and informed consent form. ARTI was defined as the presence of two or more signs of inferior respiratory tract involvement: respiratory dysfunction, characterized by fatigue, tachypnea, wheezing, crackling, and/or fever (axillary temperature of 37.8°C) on physical exam. The prodromes of URTIs were characterized by the presence of coryza, sneezing, and/or nasal obstruction. Disease history was considered in the last five days. Bronchiolitis was characterized by first episodes of wheezing in infants with URTI prodromes; children with previous episodes of pulmonary wheezing were diagnosed as

recurrent wheezing, while acute crisis was characterized as acute wheezing. Cases with crackles on pulmonary auscultation and lung consolidation on chest radiography exams were considered alveolar pneumonia, whereas cases with diffuse infiltration and with crackling on pulmonary auscultation were diagnosed as non-alveolar pneumonia.

All patients included in the study had samples of nasopharyngeal secretion collected within the first 24 h of admission. Samples were then stored on ice and sent to the Adolfo Lutz institute for Indirect Immunofluorescence (IIF), performed using a panel of five monoclonal antibodies specific for RSV, Influenza A and B, HPIV 1, 2 and 3, and Adenovirus, for the detection of seven viruses (Kit Light Diagnostic TM, Chemicon International Inc, Temecula, USA).

Variables analyzed included age, gender, admission date, clinical characteristics (cough, fever, shortness of breath, and apnea), data on physical exam (wheezing, dyspnea, cyanosis, inspiratory nasal wing collapse), clinical diagnoses, personal (cardiopathy, premature birth, and gastroesophageal reflux) and family (smoking and atopy) medical histories, patient hospital ward (general or ICU), and outcome (discharge or death).

Patients were categorized into two groups (HPIV+ or RSV+ on IIF) and subsequently compared. Medians of quantitative variables were compared using the Mann–Whitney test. Qualitative variables were analyzed by comparing proportions in each group using the Chi-square or Fisher's exact tests as applicable. The level of significance was set at 0.05. All statistical tests were performed using the Sigma Stat program version 3.5.

Results

A total of 585 patients were included in the study. HPIVs were detected in 45 patients, corresponding to 19.9% of all positive cases, and 7.7% of children overall. The RSV was the most frequently detected viral agent (26.5%). IIF was positive for influenza in 3.42% of patients and adenovirus in 1%. The clinical data of the HPIV-positive patients was compared against data from RSV-positive subjects (see Table 1). A total of 44.4% of HPIV positive patients were female versus 43.9% in the RSV-positive group. Median age was similar among HPIV+ and RSV groups, 6.8 and 7.1 months, respectively. Length of hospital stay (median six days) was also similar in the two groups. Comparing signs and symptoms of acute lower respiratory tract infection between HPIV+ and RSV+ groups revealed greater proportion of cough (96.1% vs 82.2%, $p < 0.05$) and fever (72.3% vs 48.9%, $p < 0.05$) in the RSV group. No statistically significant difference was found for the other signs and symptoms examined, including shortness of breath, wheezing, dyspnea, and cyanosis. The most frequent diagnoses in the HPIV+ group were acute wheezing (28.9%), non-alveolar pneumonia associated to acute wheezing (26.7%), and bronchiolitis (22.2%). In the RSV+ group diagnoses were bronchiolitis (35.5%), non-alveolar pneumonia associated with acute wheezing (27.1%), and acute wheezing (14.5%). Both groups were similar with regard to past personal medical history of premature birth, congenital cardiopathy, and gastroesophageal reflux disease, as well as for family history of atopy and smoking (Table 1). Of the patients in the

Table 1 – Clinical characteristics associated with infections by human parainfluenza viruses (HPIVs) and respiratory syncytial virus (RSV) (n = 585).

Clinical characteristics	PIVs (n = 45)	HRSV (n = 155)
Female gender	20 (44.4%)	68 (43.9%)
Median age (months)	6.8	7.1
Length of hospital stay (days)	6	6
Signs and symptoms of ARTI		
Cough	37 (82.2%)	149 (96.1%) ^a
Fever	22 (48.9%)	112 (72.3%) ^b
Shortness of breath	28 (62.2%)	115 (74.2%)
Wheezing	30 (66.7%)	85 (54.8%)
Dyspnea	27 (60%)	96 (61.9%)
Cyanosis	4 (8.9%)	4 (2.58%)
ARTI diagnoses		
Acute wheezing	13 (28.9%)	23 (14.5%)
Non-alveolar pneumonia	12 (26.7%)	42 (27.1%)
Bronchiolitis	10 (22.2%)	55 (35.5%)
Personal medical history		
Premature birth	6 (13.3%)	13 (8.4%)
Congenital cardiopathy	4 (8.9%)	8 (5.2%)
Gastroesophageal reflux	5 (11.1%)	11 (7.1%)
Family medical history		
Atopy	20 (44.4%)	56 (36.1%)
Smoking	6 (13.3%)	20 (12.9%)
ICU admission	7 (15.5%)	17 (10.9%)
Death	4 (8.9%)	2 (1.3%) ^c

^a p = 0.0037.

^b p = 0.0058.

^c p = 0.024.

HRSV+ group, 10.9% were ICU cases compared with 15.5% in the HPIV+ group (Table 1). A greater number of deaths were recorded in the HPIV+ group compared with the HRSV+ group (8.9% vs. 1.29%, $p < 0.05$) (Table 1).

Of the HPIVs detected, the most frequent type was HPIV-3 (84.4%) followed by HPIV-2 (8.89%) and HPIV-1 (6.67%).

The peak detection rate of the HRSV occurred in the autumn season during the study period. In the first two years of the study, a higher incidence of HPIVs was observed accompanying the HRSV peaks (Fig. 1).

Discussion

The HPIVs are important etiologic agents of acute respiratory infection in childhood and rank second among the most common diseases in this age group by some authors. HPIVs predominantly cause lower respiratory tract infection in infants and young children,¹²⁻¹⁴ and the significance of the virus has been underappreciated.⁹

In the present investigation, the prevalence of HPIVs was 7.7%, the second most common virus after HRSV, whose prevalence was 26.5%. This result corroborates the findings of other studies with similar design.¹⁵⁻¹⁷ We adopted IIF as the method of viral investigation, as it is a reliable technique with 50–75% sensitivity and excellent specificity.^{12,18} Some authors claim that molecular methods (PCR and Real-Time PCR) offer 10–20%

greater sensitivity for detecting HPIVs over culture and IIF techniques,^{19,20} but with higher cost.

Earlier studies have reported higher prevalence of HPIVs than in the present study but had different study designs and viral detection methods. For instance, the study by Rihkanen et al., investigating the etiology of croup in patients attending emergency services, found HPIVs, particularly type 1, in 29.1% of patients.²¹ HPIV1 is known to be the main etiologic agent of croup, accounting for 56–74% of such cases. A cohort study conducted in Spain, involving patients hospitalized for ARTI aged up to 14 years using PCR as the viral detection method, showed a 9.9% HPIV prevalence, ranking third among the most frequent viruses after the HRSV, Rhinovirus and Adenovirus.⁶ In a study aiming to determine the etiology of community-acquired pneumonia in hospitalized patients aged younger than five in Salvador city in northeastern Brazil, HPIVs ranked second among the most prevalent agents, followed by Rhinovirus.²²

The prevalence of HPIV-3 was substantially higher than HPIV-1 and 2 in this study, whereas HPIV-4 was not studied. It is known that HPIV-3 is the HPIV type most frequently detected in hospitalized patients, particularly among young infants, and associated with the diagnoses of bronchiolitis and pneumonia.^{12,23}

Other studies have shown HPIV-3 to be the most prevalent in hospitalized patients, such as the work by Calvo et al.⁶ in addition to studies conducted in Brazil by Fe et al.²⁴ assessing the prevalence of HPIVs in patients up to 16 years old with ARTI, and the study by Veja-Briceño et al. conducted in Chile.²⁵ Evidence confirms that HPIV-1 is associated with the diagnosis of croup, but not a single case of croup or laryngitis associated to croup was found in our study, perhaps because our cases were often less severe, not requiring hospitalization.

Comparing the clinical data of HPIVs and HRSV infected groups, we found no statistically significant differences in terms of hospital stay or need for ICU admission. With regard to signs and symptoms of ARTI, there were greater proportions of cough and fever in the HRSV-infected group. Despite this association, because signs and symptoms were very unspecific and general, none of these could be attributed to a specific agent.^{25,26}

The highest rates of HPIV isolation (detection) occurred during autumn in the first two years of the study, coinciding with peak incidence of the HRSV. The classic seasonal pattern of HPIV infection is described in the United States, with biannual peaks of HPIV1 during fall and winter, and annual peaks of HPIV3, particularly at the end of winter or in spring.^{9,27} Studies in tropical countries are scarce, although one investigation reported seasonal peaks of HPIV3 at the end of winter and in spring in Rio de Janeiro,²⁸ while other studies carried out in Northeastern and Southern Brazil both showed peaks during October, coinciding with spring.^{24,29}

Evidence shows that HPIV infections tend to be less severe than those caused by HRSV.⁶ More deaths occurred in children hospitalized for HPIVs than among HRSV-infected patients. The deaths occurred in children outside the age group at higher risk for developing severe disease (less than three months old), but all these cases had contributing medical factors, with two having history of premature birth, one recurrent wheezing case with gastroesophageal reflux, and another




Fig. 1 – Distribution of cases by month and number of patients with human parainfluenza viruses (HPIV) or with human respiratory syncytial virus (VSR).

with delayed neuro-psychomotor development secondary to neonatal anoxia and recurrent wheezing. The prevention strategy of administering immunoglobulin is well established for HRSV, and studies are underway for vaccines against this agent. For HPIVs however, viruses with significant impact in terms of morbidity and mortality, no established strategy is currently in place.

In conclusion, the impact caused by HPIVs infections is significant in hospitalized children, where this agent should be considered as a cause of ARTI. Longitudinal studies determining the prevalence of respiratory viruses should be encouraged to help devise prevention and treatment strategies.

Funding

This study was funded by FAPESP – Fundação de Amparo à Pesquisa do Estado de São Paulo.

Conflicts of interest

The authors declare no conflicts of interest.

REFERENCES

- UNICEF. World Health Organization (WHO). Pneumonia: the forgotten Killer of children. Geneva: UNICEF/WHO; 2006, 40 pp.
- Williams BG, Gouws E, Boschi-Pinto C, Bryce J, Dye C. Estimates of world-wide distribution of child deaths from acute respiratory infections. Lancet Infect Dis. 2002;2:25–32.
- Lourenço LG, Júnior JBS, Rahal P, Souza FP, Zanetta DMT. Infecção pelo Vírus Sincicial Respiratório em Crianças. Pulmão RJ. 2005;14:59–68.
- Pilger DA, Cantarelli VV, Amantea SL, et al. Detection of human bocavirus and human metapneumovirus by real-time PCR from patients with respiratory symptoms in Southern Brazil. Mem Inst Oswaldo Cruz. 2011;106:56–60.
- Bonzel L, Tenenbaum T, Schrotten H, Schildgen O, Schweitzer-Krantz S, Adams O. Frequent detection of viral coinfection in children hospitalized with acute respiratory tract infection using a real-time polymerase chain reaction. Pediatr Infect Dis J. 2008;27:589–94.
- Calvo C, Garcia-Garcia ML, Ambrona P, et al. The burden of infections by parainfluenza virus in hospitalized children in Spain. Pediatr Infect Dis J. 2011;30:792–4.
- Chiu SS, Chan KH, Chen H, et al. Virologically confirmed population-based burden of hospitalization caused by respiratory syncytial virus, adenovirus, and parainfluenza viruses in children in Hong Kong. Pediatr Infect Dis J. 2010;29:1088–92.
- Yang HT, Jiang Q, Zhou X, et al. Identification of a natural human serotype 3 parainfluenza virus. Virol J. 2011;8:58.
- Weinberg GA. Parainfluenza viruses: an underappreciated cause of pediatric respiratory morbidity. Pediatr Infect Dis J. 2006;25:447–8.
- Lee MS, Mendelman PM, Sangli C, Cho I, Mathie SL, August MJ. Half-life of human parainfluenza virus type 3 (HPIV3) maternal antibody and cumulative proportion of HPIV3 infection in young infants. J Infect Dis. 2001;183:1281–4.
- Wolf DG, Greenberg D, Kalkstein D, et al. Comparison of human metapneumovirus, respiratory syncytial virus and influenza A virus lower respiratory tract infections in

- hospitalized young children. *Pediatr Infect Dis J.* 2006;25:320-4.
12. Henrickson KJ. Parainfluenza viruses. *Clin Microbiol Rev.* 2003;16:242-64.
 13. Sato M, Wright PF. Current status of vaccines for parainfluenza virus infections. *Pediatr Infect Dis J.* 2008;27:S123-5.
 14. Teo WY, Rajadurai VS, Sriram B. Morbidity of parainfluenza 3 outbreak in preterm infants in a neonatal unit. *Ann Acad Med Singapore.* 2010;39:837-46.
 15. Iwane MK, Edwards KM, Szilagyi PG, et al. Population-based surveillance for hospitalizations associated with respiratory syncytial virus, influenza virus, and parainfluenza viruses among young children. *Pediatrics.* 2004;113:1758-64.
 16. Noyola DE, Arteaga-Dominguez G. Contribution of respiratory syncytial virus, influenza and parainfluenza viruses to acute respiratory infections in San Luis Potosi, Mexico. *Pediatr Infect Dis J.* 2005;24:1049-52.
 17. Pecchini R, Berezin EN, Felicio MC, et al. Incidence and clinical characteristics of the infection by the respiratory syncytial virus in children admitted in Santa Casa de Sao Paulo Hospital. *Braz J Infect Dis.* 2008;12:476-9.
 18. Greenhow TL, Weinrub PS. Utility of direct fluorescent antibody testing of nasopharyngeal washes in children with and without respiratory tract illness. *Pediatr Infect Dis J.* 2006;25:502-6.
 19. Freymuth F, Vabret A, Cuvillon-Nimal D, et al. Comparison of multiplex PCR assays and conventional techniques for the diagnostic of respiratory virus infections in children admitted to hospital with an acute respiratory illness. *J Med Virol.* 2006;78:1498-504.
 20. Syrmis MW, Whiley DM, Thomas M, et al. A sensitive, specific, and cost-effective multiplex reverse transcriptase-PCR assay for the detection of seven common respiratory viruses in respiratory samples. *J Mol Diagn.* 2004;6:125-31.
 21. Peltola V, Heikkinen T, Ruuskanen O. Clinical courses of croup caused by influenza and parainfluenza viruses. *Pediatr Infect Dis J.* 2002;21:76-8.
 22. Nascimento-Carvalho CM, Ribeiro CT, Cardoso MR, et al. The role of respiratory viral infections among children hospitalized for community-acquired pneumonia in a developing country. *Pediatr Infect Dis J.* 2008;27:939-41.
 23. Gomez M, Mufson MA, Dubovsky F, Knightly C, Zeng W, Losonsky G. Phase-I study MEDI-534, of a live, attenuated intranasal vaccine against respiratory syncytial virus and parainfluenza-3 virus in seropositive children. *Pediatr Infect Dis J.* 2009;28:655-8.
 24. Fe MM, Monteiro AJ, Moura FE. Parainfluenza virus infections in a tropical city: clinical and epidemiological aspects. *Braz J Infect Dis.* 2008;12:192-7.
 25. Vega-Briceno LE, Pulgar BD, Potin SM, Ferres GM, Sanchez DI. Clinical and epidemiological manifestations of parainfluenza infection in hospitalized children. *Rev Chilena Infectol.* 2007;24:377-83.
 26. Thomazelli LM, Vieira S, Leal AL, et al. Surveillance of eight respiratory viruses in clinical samples of pediatric patients in southeast Brazil. *J Pediatr.* 2007;83:422-8.
 27. Hall CB. Respiratory syncytial virus and parainfluenza virus. *N Engl J Med.* 2001;344:1917-28.
 28. Nascimento JP, Siqueira MM, Sutmoller F, et al. Longitudinal study of acute respiratory diseases in Rio de Janeiro: occurrence of respiratory viruses during four consecutive years. *Rev Inst Med Trop Sao Paulo.* 1991;33:287-96.
 29. Straliotto SM, Siqueira MM, Muller RL, Fischer GB, Cunha ML, Nestor SM. Viral etiology of acute respiratory infections among children in Porto Alegre, RS, Brazil. *Rev Soc Bras Med Trop.* 2002;35:283-91.