



The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



Original Article

Knowledge about clinical presentation, prevention strategies and sexual transmission of Zika virus infection among women of reproductive age in an endemic area



Erika R. Manuli ^{a,b}, Geovana M. Pereira ^b, Maria Cristina Bernat ^c,
Celia R. Novaes ^d, Ester C. Sabino ^{a,b}, Vivian I. Avelino-Silva ^{a,b,*}

^a Department of Infectious and Parasitic Diseases, Faculdade de Medicina da Universidade de São Paulo, São Paulo, SP, Brazil

^b Instituto de Medicina Tropical, Faculdade de Medicina da Universidade de São Paulo, São Paulo, SP, Brazil

^c Centro de Saude Escola Geraldo de Paula Souza, Faculdade de Saúde Pública, Universidade de São Paulo, São Paulo, SP, Brazil

^d Centro de Saude I "Dr. Victor Araujo Homem de Mello", Departamento de Gerenciamento Ambulatorial da Capital, Secretaria de Estado da Saúde, São Paulo, São Paulo, SP, Brazil

ARTICLE INFO

Article history:

Received 1 May 2021

Accepted 20 September 2021

Available online 8 October 2021

Keywords:

Health knowledge
Attitudes and practice
Zika virus
Pregnancy
Sexual transmission
Women of childbearing age

ABSTRACT

Background: The recognition of the causal association between Zika virus (ZIKV) infection during pregnancy and congenital abnormalities including microcephaly underlines the importance of preventing this disease in pregnant women (PW) and women of childbearing age (WCA). Although Brazil and other Latin American countries reported a significant reduction in the number of ZIKV infections in recent years, epidemic waves can recur in settings with previous outbreaks as conditions for transmission remain optimal and susceptible populations are continuously replenished.

Methods: In this cross-sectional study, we enrolled 64 PW and 260 non-pregnant WCA attending routine medical appointments in two primary care units in Sao Paulo, Brazil, and assessed knowledge and attitudes about ZIKV infection and prevention.

Results: Most women reported knowing that ZIKV is transmitted through the bite of *Aedes* mosquitoes, and most knew that acute symptoms are similar to those seen in Dengue infection. Furthermore, most participants correctly described that ZIKV infection during pregnancy may cause detrimental outcomes for the newborn. However, most ignored that ZIKV infection can be asymptomatic, and only 15% knew about the risk of ZIKV sexual transmission. We found no statistically significant differences between PW and WCA regarding knowledge about ZIKV sexual transmission. Knowledge about ZIKV sexual transmission was significantly associated with education; among participants with ≤ 12 schooling years, only 9.0% (95%CI 3.4-18.5%) correctly answered that ZIKV can be sexually transmitted, compared to 12.9% (95%CI 8.2-18.8%) among participants with 12-14 schooling years, and to 24.4% (95%CI 15.9-34.9%) of participants with ≥ 15 schooling years ($p = 0.015$). Education remained independently associated with knowledge about sexual transmission of ZIKV in

* Corresponding author.

E-mail address: viviansilva87@gmail.com (V.I. Avelino-Silva).

<https://doi.org/10.1016/j.bjid.2021.101629>

1413-8670/© 2021 Sociedade Brasileira de Infectologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

a multivariate logistic regression model adjusted for age, race and pregnancy status ($p = 0.022$).

Conclusion: Our findings underscore the urgent need of educational and family planning programs that may help prevent detrimental outcomes of ZIKV infection in an endemic area of Brazil.

© 2021 Sociedade Brasileira de Infectologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Introduction

Since 2016, Brazilian and international scientific groups have been dedicating intense efforts to understand Zika virus (ZIKV) infection.¹ This interest has been raised following a massive epidemic wave in Latin America and the detection of a significant increase in the number of newborns with microcephaly and other congenital malformations causally associated with ZIKV infection during pregnancy. Brazil was the first country to identify an increase in microcephaly cases, and the causal association with ZIKV infection was supported by studies showing that the virus was present in damaged fetal neural tissues; moreover, ZIKV was shown to have a neurotoxic effect in both *in vitro* and animal models.²⁻⁴

As seen with other viruses belonging to the Flavivirus family, ZIKV is transmitted through the bite of *Aedes* mosquitos, and *Aedes aegypti* seems to be the main vector for the disease in the Americas.⁵ Blood transfusions can also be a potential source of ZIKV infection.^{6, 7} Sexual intercourse is another non-vectorial mode of ZIKV transmission, particularly from a male infected partner^{8, 9}; viral RNA has been shown to persist in the semen of a few patients for several months after symptoms onset, with potential transmissibility lasting 6-7 weeks.¹⁰

The first Brazilian cases of ZIKV infection were detected in 2015 in the Northeast region of the country, with subsequent expansion to states in the Central-West and Southeast regions.² In the state of Sao Paulo, ZIKV outbreaks were more significant in rural and countryside cities and in coastal cities.¹¹ After the peak case-detection in 2016, Brazil and other Latin American countries reported an important reduction in the number of ZIKV infections.¹² However, new epidemic waves can recur in settings with previous outbreaks as conditions for transmission remain optimal and susceptible populations are continuously replenished. Furthermore, the World Health Organization highlighted in a recent report that approximately 61 countries with no reports of ZIKV occurrence have local conditions for the spread of the virus, with naturally occurring competent vectors, ideal climate settings and vulnerable populations. This report also discusses the importance of providing family planning strategies and travel orientation in countries with ZIKV occurrence.¹³ Since ZIKV still lacks effective antiviral treatments or vaccines, health services and public health institutions should be prepared to respond to new disease outbreaks.¹⁴ One essential way of being prepared is supporting vulnerable populations such as women of childbearing age (WCA) with knowledge about the disease and how to prevent it.¹⁵

In this study we enrolled pregnant women (PW) and non-pregnant WCA in two primary care units in Sao Paulo, Brazil, and assessed knowledge and attitudes about ZIKV infection and prevention. We also explored if the recent ZIKV outbreak had an impact on family planning issues in this population.

Methods

Study design, population and setting

In this cross-sectional study, we enrolled PW and non-pregnant WCA attending routine medical appointments at two primary care units, both in central areas of Sao Paulo city, Brazil. Consenting women aged 18-45 years old living in Sao Paulo city were considered eligible; we excluded women with current or past ZIKV infection and those who had infertility due to any cause.

Study procedures

After obtaining informed consent for participation from each woman, a trained study investigator administered the study questionnaire that included questions on knowledge about ZIKV infection, knowledge and attitudes towards prevention strategies, and the impact of the recent ZIKV outbreak on family planning. Pregnancy status at the time of interview was self-reported. Responses were registered in a standardized electronic form using REDCap platform.¹⁶

Ethical aspects

All participants provided informed consent upon inclusion in the study. We collected no participant-identifiable information, and all study forms were kept in locked cabinets or password-secured electronic folders in accordance with Good Clinical Practice recommendations. The study has been approved by the Ethics committee at University of Sao Paulo Medical School (approval reference number 2.358.114) as well as local ethics committees at each primary care unit. As a benefit from participation, the study investigator offered to review all questionnaire answers and clarify any remaining uncertainties; a printed booklet produced by the Brazilian Ministry of Health containing information on ZIKV was also offered to all participants at the end of study participation.¹⁷

Statistical analysis

Data was analyzed using descriptive statistics. Comparisons between PW and WCA were preformed using chi-squared tests or Fisher's exact test for categorical variables, and t-tests or Wilcoxon Rank-sum tests for numerical variables, as appropriate. A multivariate logistic regression model was used to explore independent predictors of knowledge about ZIKV sexual transmission. We used Stata 15.1 (StataCorp. College Station, TX: StataCorp LP), with a two-tailed significance level of 0.05 for all analyses.

Results

Between December 2017 and July 2018, 409 PW and WCA were invited to participate in the study; of those, 85 were excluded (21 refused or withdrew participation; 4 lived in other cities; 37 were older than 45 or younger than 18 years old and 23 were excluded due to infertility related to past/current ZIKV infection). Among 324 women included in the study, 64 (20%) were pregnant. Demographic characteristics of study participants, overall and according to pregnancy status, are presented in Table 1. PW and non-pregnant WCA were similar regarding age, race/ethnicity, income, and education.

Out of 64 PW enrolled in the study, 22 (34%) reported they had planned to get pregnant; 21 (33%) said they were neither planning nor avoiding pregnancy; and 21 (33%) said they got pregnant despite taking precautions to avoid pregnancy. Only three out of 12 women who reported trying to postpone pregnancy attributed it to fear of ZIKV infection (Table 2).

Table 3 presents participants' knowledge and attitudes towards ZIKV infection. Most women reported knowing that ZIKV is transmitted through the bite of *Aedes* mosquitos, and

Table 2 – Family planning aspects among 64 pregnant women enrolled in the study.

How did this pregnancy happen? (%)	
I planned to get pregnant	22 (34%)
I did not plan, di did not avoid getting pregnant	21 (33%)
I was taking precautions to avoid pregnancy	21 (33%)
Did you try to avoid or postpone pregnancy? (%)	
No	52 (81%)
Yes	12 (19%)
Did you try to avoid or postpone pregnancy because of fear of ZIKV infection? (%)	
No	9 (75%)
Yes	3 (25%)

ZIKV: Zika virus

most knew that acute symptoms are similar to those seen in Dengue infection. Furthermore, most participants correctly described that ZIKV infection during pregnancy may cause detrimental outcomes for the newborn. However, most ignored that ZIKV infection can be asymptomatic, and only 49 (15%) knew about the risk of ZIKV sexual transmission. Surprisingly, we found no statistically significant differences between PW and WCA in the knowledge and attitudes towards ZIKV infection, except for higher knowledge about ZIKV cases in the city ($p = 0.047$) and lower condom use by the sexual partner ($p = 0.001$) among PW.

Predictors of knowledge about ZIKV sexual transmission

Given the strikingly low percentage of knowledge about ZIKV sexual transmission among women of childbearing age regardless of pregnancy status, we further explored if age category, race, education and income were associated with this finding. We found a statistically significant association

Table 1 – Demographic characteristics of study participants, overall and according to pregnancy status.

Characteristics	Total N = 324	Pregnant women N = 64	Non-pregnant women of childbearing age N = 260	p-value
Age	33 (27-39)	32 (26-37)	34 (27-40)	0.087
Race/ethnicity (%)				0.327
White/Caucasian	115 (35)	22 (34)	93 (36)	
Black	50 (15)	10 (16)	40 (15)	
Mixed	129 (40)	24 (38)	105 (40)	
Asian	22 (7)	4 (6)	18 (7)	
Native	2 (1)	1 (2)	1 (<1)	
Other/not informed	6 (2)	3 (5)	3 (1)	
Household income in US dollars (%)				0.166
<100	8 (2)	3 (5)	5 (2)	
100-200	36 (11)	5 (8)	31 (12)	
200-300	73 (23)	9 (14)	64 (25)	
300-400	83 (26)	15 (23)	68 (26)	
400-600	55 (17)	14 (22)	41 (16)	
>600	66 (20)	17 (27)	49 (19)	
Not informed	3 (1)	1 (2)	2 (1)	
Schooling years	12 (12-15)	12 (12-15)	12 (12-15)	0.703
Number of children	1 (0-2)	1 (0-1)	1 (0-2)	0.096
Primary care unit (%)				-
Unit 1	171 (53)	57 (89)	114 (44)	
Unit 2	153 (47)	7 (11)	146 (56)	

Numeric variables are presented as medians and interquartile ranges

Table 3 – Knowledge and attitudes concerning ZIKV infection among study participants, overall and according to pregnancy status.

Item / N (%) of correct responses	Total N = 324	Pregnant women N = 64	Non-pregnant women of childbearing age N = 260	p-value
Reported existing ZIKV cases in the city	186 (58)	44 (69)	142 (55)	0.047
ZIKV can be transmitted through the bite of <i>Aedes</i>	290 (90)	58 (91)	232 (89)	0.744
The infected person almost always has symptoms	237 (73)	45 (70)	192 (74)	0.511
ZIKV can be transmitted through sexual intercourse	49 (15)	13 (20)	36 (14)	0.196
ZIKV symptoms and dengue symptoms are similar	295 (92)	57 (90)	238 (92)	0.716
Infection with ZIKV during pregnancy can cause severe problems to the infant	319 (98)	63 (98)	256 (98)	1.000
Microcephaly is the only abnormality caused by ZIKV infection during pregnancy	116 (36)	21 (33)	95 (37)	0.578
If the PW is infected towards the end of pregnancy, there are no risks for the infant	198 (61)	36 (56)	162 (63)	0.336
If a pregnant woman get ZIKV, she can be treated and eliminate the virus	97 (30)	14 (22)	83 (32)	0.116
I avoid going to places infested with mosquitos:				
Always	92 (28)	22 (34)	70 (27)	0.195
Almost always	81 (25)	16 (25)	65 (25)	
I neither avoid nor attend unnecessarily	58 (18)	15 (23)	43 (17)	
Almost never avoid	34 (10)	4 (6)	30 (12)	
Never avoid	59 (18)	7 (11)	52 (20)	
I use mosquito repellents:				
Always	93 (29)	18 (28)	75 (29)	0.478
Almost always	45 (14)	12 (19)	33 (13)	
Sometimes	88 (27)	16 (25)	72 (28)	
Almost never	30 (9)	8 (13)	22 (8)	
Never	68 (21)	10 (16)	58 (22)	
I wear long clothes to cover my skin:				
Always	51 (16)	17 (27)	34 (13)	0.116
Almost always	47 (15)	7 (11)	40 (15)	
Sometimes	81 (25)	15 (23)	66 (25)	
Almost never	39 (12)	6 (9)	33 (13)	
Never	106 (33)	19 (30)	87 (33)	
My partner uses mosquito repellents:				
Always	41 (15)	6 (10)	35 (17)	0.289
Almost always	30 (11)	5 (8)	25 (12)	
Sometimes	44 (16)	8 (13)	36 (17)	
Almost never	22 (8)	6 (10)	16 (8)	
Never	132 (49)	37 (60)	95 (46)	
My partner wears condoms:				
Always	73 (27)	4 (6)	69 (33)	0.001
Almost always	29 (11)	7 (11)	22 (11)	
Sometimes	27 (10)	6 (10)	21 (10)	
Almost never	13 (5)	5 (8)	8 (4)	
Never	130 (48)	41 (65)	89 (43)	

between education and knowledge about ZIKV sexual transmission; among participants with ≤ 12 schooling years, only 9% (95% CI 3-19%) correctly answered that ZIKV can be sexually transmitted; this percentage was 13% (95% CI 8-19%) among participants with 12-14 schooling years, and 24% (95% CI 16-35%) among participants with ≥ 15 schooling years ($p = 0.015$; Table 4). Age category, race, and income were not significantly associated with knowledge about ZIKV sexual transmission in univariate analysis (Table 4). In a multivariate logistic regression model adjusted for age, race and pregnancy status, education remained independently associated with knowledge about sexual transmission of ZIKV; participants with > 12 schooling years had 2.10 times the odds of

reporting this knowledge compared to those with ≤ 12 schooling years (95% CI 1.11-3.95; $p = 0.022$; Table 5).

Discussion

In this cross-sectional study, we addressed knowledge and attitudes about ZIKV infection among PW and non-pregnant WCA. We showed that the knowledge about transmission by mosquitos and the potential to cause congenital malformations when infection occurs during pregnancy was generally high. However, the knowledge about sexual transmission was very low among both PW and non-pregnant WCA. These

Table 4 – Factors associated with knowledge about ZIKV sexual transmission in univariate analysis.

	Knows about ZIKV sexual transmission N = 49	Doesn't know about ZIKV sexual transmission N = 275	Univariate analysis p-value
Pregnancy status (%)			
Pregnant	13 (20)	51 (80)	0.196
Non-pregnant	36 (14)	224 (86)	
Age (%)			
<27 years old	11 (14)	68 (86)	0.522
27-32 years old	13 (18)	59 (82)	
33-38 years old	16 (18)	74 (82)	
≥39 years old	9 (11)	74 (89)	
Race (%)			
White/Caucasian/Asian	20 (15)	117 (85)	0.806
Black/Mixed/Other	29 (16)	157 (84)	
Education (%)			
<12 years	6 (9)	61 (91)	0.015
12-14 years	22 (13)	149 (87)	
≥15 years	21 (24)	65 (76)	
Income (%)			
<200 US dollars	7 (16)	37 (84)	0.153
200-400 US dollars	17 (11)	139 (89)	
400-600 US dollars	9 (16)	46 (84)	
>600 US dollars	15 (23)	51 (77)	

findings are alarming, particularly in the context of the recent outbreak of ZIKV in Brazil, with ample repercussion in mass communication vehicles. Furthermore, during study design our group had initially hypothesized that PW would have higher levels of knowledge and better prevention attitudes compared to non-pregnant WCA, given multiple opportunities of learning during antenatal care consultations.¹⁸ We were surprised to see that PW and WCA had similar knowledge and prevention attitudes, except for the knowledge about ZIKV cases in the city and use of condoms by the sexual partner. This was expected, since contraception is certainly one of the main reasons for condom use among non-pregnant couples and will be less frequently adopted if the woman is already pregnant. However, it also emphasizes that prevention of ZIKV sexual transmission to PW may face significant challenges in endemic countries. Knowledge about ZIKV sexual transmission was reported by only 15% of participants in our study and was similarly low among PW and non-pregnant

WCA. Education was significantly associated with knowledge about ZIKV sexual transmission in both univariate and multivariate analysis. Finally, we also found that a very small number of PW reported they had postponed pregnancy due to the ZIKV epidemic in our setting.

A few prior studies have explored knowledge, attitudes and practices of WCA towards ZIKV infection. Knowledge about ZIKV sexual transmission was assessed among PW and women attending antenatal care clinics, and was shown to be lacking in the majority of participants in a study conducted in the USA¹⁹; similarly, 74% of respondents in a study conducted in Trinidad and Tobago²⁰ and 63% in a study conducted in Greece²¹ ignored that ZIKV could be transmitted through sexual intercourse. In a survey including male and female participants in the USA, only 27% identified abstinence or using condoms during intercourse as precautions for ZIKV infection in PW traveling to endemic areas.¹⁸ In a study published by Borges et al., who interviewed 526 women aged 18 to 49 years

Table 5 – Factors associated with knowledge about ZIKV sexual transmission in multivariate analysis.

	Odds ratio	95% confidence interval	Multivariate analysis p-value
Pregnancy status			
Pregnant	1.47	0.72-3.00	0.296
Non-pregnant	1 (referent)	-	
Age			
<27 years old	1 (referent)	-	0.577
27-32 years old	1.29	0.53-3.12	
33-38 years old	1.29	0.55-3.00	
≥39 years old	0.83	0.32-2.15	
Race			
White/Caucasian/Asian	0.85	0.45-1.60	0.608
Black/Mixed/Other	1 (referent)	-	
Education			
≤12 years	1 (referent)	-	0.022
>12 years	2.10	1.11-3.95	

old in the Northeast of Brazil, awareness about the congenital syndrome associated with ZIKV infection during pregnancy was high, but only 50% of participants knew about the risk of sexual transmission. The higher proportion of women with knowledge about ZIKV sexual transmission compared with our sample may be partially explained by the fact that the Northeast of the country was the epicenter of the disease, and the study was conducted few months after the peak of ZIKV epidemic.²² However, this proportion is still not ideal, and further studies should be conducted to assess if this knowledge is retained years after the peak of the epidemic. In accordance with our findings, education and other social determinants have been associated with knowledge and attitudes about ZIKV in previous publications,^{21, 23} suggesting that WCA in more vulnerable settings should be prioritized in public health policies and prevention strategies. Although vector transmission is undoubtedly the main route of infection for ZIKV, spread through sexual contact also contributes to the transmission chain²⁴ and should be prevented particularly among pregnant women. Recommendations for abstinence and safer sex in high ZIKV transmission areas have been issued by Brazilian²⁵ as well as international organizations.^{26, 27}

Our study has a few limitations. We restricted selection of participants to a convenience sample in two primary care units, both located in central, more affluent areas of Sao Paulo city. Therefore, our study sample comprised women with higher socioeconomic status than the average population in our setting. Women with no access to healthcare were not available for inclusion in our study. These limitations emphasize that the overall knowledge about ZIKV is probably even lower, reinforcing the urgent need for educational actions that may help prevent ZIKV detrimental outcomes.

Our study included PW and non-pregnant WCA between December 2017 and July 2018, over a year after the peak detection of ZIKV cases in the country. The improvements in epidemiological figures may have decreased the pressure for public health interventions supporting ZIKV prevention, particularly concerning PW and non-pregnant WCA. However, Brazil has optimal vector and climate conditions for a recurring outbreak, and actions supporting education and self-care should be intensified. Other countries with potential risk of future ZIKV outbreaks should also implement such strategies in order to avoid subsequent preventable outbreaks of congenital abnormalities.¹³

Conflicts of interest

None to declare.

REFERENCES

- Fauci AS, Morens DM. Zika virus in the Americas—yet another arbovirus threat. *N Engl J Med*. 2016;374:601–4.
- Lowe R, Barcellos C, Brasil P, et al. The Zika virus epidemic in Brazil: from discovery to future implications. *Int J Environ Res Public Health*. 2018;15(1):96.
- Schuler-Faccini L, Ribeiro EM, Feitosa IM, et al. Possible association between Zika virus infection and microcephaly - Brazil, 2015. *MMWR Morb Mortal Wkly Rep*. 2016;65:59–62.
- Teixeira MG, Costa Mda C, de Oliveira WK, Nunes ML, Rodrigues LC. The epidemic of Zika virus-related microcephaly in Brazil: detection, control, etiology, and future scenarios. *Am J Public Health*. 2016;106:601–5.
- Domingues CMAS, Teixeira AMS. Coberturas vacinais e doenças imunopreveníveis no Brasil no período 1982-2012: avanços e desafios do Programa Nacional de Imunizações. *Epidemiol Serv Saúde*. 2013;22:9–27.
- Basile K, Kok J, Dwyer DE. Zika virus: what, where from and where to? *Pathology*. 2017;49:698–706.
- Petersen E, Wilson ME, Touch S, et al. Rapid spread of Zika virus in the Americas—implications for public health preparedness for mass gatherings at the 2016 Brazil olympic games. *Int J Infect Dis*. 2016;44:11–5.
- Sherley M, Ong CW. Sexual transmission of Zika virus: a literature review. *Sex Health*. 2018;15(3):183–99.
- Counotte MJ, Kim CR, Wang J, et al. Sexual transmission of Zika virus and other flaviviruses: a living systematic review. *PLoS Med*. 2018;15:e1002611.
- Musso D, Gubler DJ. Zika Virus. *Clin Microbiol Rev*. 2016;29:487–524.
- Governo do Estado de Sao Paulo. Secretaria de Estado da Saude. Centro de Vigilancia Epidemiologica “Prof. Alexandre Vranjac”. Dados estatísticos. Available at: <https://www.saude.sp.gov.br/cve-centro-de-vigilancia-epidemiologica-prof.-alexandre-vranjac/areas-de-vigilancia/doencas-de-transmissao-por-vetores-e-zoonoses/arboviroses-urbanas/zika-virus/dados-estatisticos>. Access date: October 8th, 2021.
- WHO Region of the Americas/Pan American Health Organization. PLISA Health Information Platform for the Americas: Cases of Zika virus disease, by country or territory. Available at: <https://www3.paho.org/data/index.php/en/mnu-topics/zika/524-zika-weekly-en.html>. Accessed October 8th 2021.
- World Health Organization. Global Overview ZIKA Epidemiology Update. Available at: <https://www.who.int/emergencies/diseases/zika/zika-epidemiology-update-july-2019.pdf?ua=1>. Accessed October 1 2020.
- Pielnaa P, Al-Saadawe M, Saro A, et al. Zika virus-spread, epidemiology, genome, transmission cycle, clinical manifestation, associated challenges, vaccine and antiviral drug development. *Virology*. 2020;543:34–42.
- Centers for Disease Control and Prevention. Zika Virus Prevention and Transmission. Available at: <https://www.cdc.gov/zika/prevention/index.html>. Accessed October 8th, 2021.
- Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform*. 2009;42:377–81.
- Brasil. Ministério da Saúde. Vírus ZIKA Informações ao Público. Available at: http://biblioteca.cofen.gov.br/wp-content/uploads/2016/01/Cartilha_Zika_MS.pdf. Accessed October 8th, 2021.
- Samuel G, DiBartolo-Cordovano R, Taj I, et al. A survey of the knowledge, attitudes and practices on Zika virus in new York City. *BMC Public Health*. 2018;18:98.
- Berenson AB, Trinh HN, Hirth JM, Guo F, Fuchs EL, Weaver SC. Knowledge and prevention practices among U.S. pregnant immigrants from Zika virus outbreak areas. *Am J Trop Med Hyg*. 2017;97:155–62.
- Pooransingh S, Parasram R, Nandram N, Bhagwandeem B, Dialsingh I. Zika virus disease-knowledge, attitudes and practices among pregnant women-implications for public health practice. *Public Health*. 2018;165:146–51.
- Mouchtouri VA, Papagiannis D, Katsioulis A, Rachiotis G, Dafopoulos K, Hadjichristodoulou C. Knowledge, attitudes, and practices about the prevention of mosquito bites and Zika

- virus disease in pregnant women in greece. *Int J Environ Res Public Health*. 2017;14.
22. Borges ALV, Moreau C, Burke A, Dos Santos OA, Chofakian CB. Women's reproductive health knowledge, attitudes and practices in relation to the Zika virus outbreak in northeast Brazil. *PLoS One*. 2018;13:e0190024.
 23. Maharajan MK, Rajiah K, Belotindos JS, Basa MS. Social determinants predicting the knowledge, attitudes, and practices of women toward Zika virus infection. *Front Public Health*. 2020;8:170.
 24. Denes A, Ibrahim MA, Oluoch L, Tekeli M, Tekeli T. Impact of weather seasonality and sexual transmission on the spread of Zika fever. *Sci Rep*. 2019;9:17055.
 25. Duarte G, Miranda AE, Bermudez XPD, Saraceni V, Martinez-Espinosa FE. Brazilian protocol for sexually transmitted infections 2020: Zika virus infection. *Rev Soc Bras Med Trop*. 2021;54(suppl 1):e2020609.
 26. Polen KD, Gilboa SM, Hills S, et al. Update: interim guidance for preconception counseling and prevention of sexual transmission of Zika virus for men with possible Zika virus exposure - United States, August 2018. *MMWR Morb Mortal Wkly Rep*. 2018;67:868-71.
 27. World Health Organization. WHO guidelines for the prevention of sexual transmission of Zika virus. Available at: <https://www.who.int/publications/i/item/9789241550482>. Access date: October 8th, 2021.